

Appendix F: List of Evidence Identified from Literature Search

(Cross reference to Section 3.3 in Report)

Highlighted reference was relevant to Regional Project

Highlighted reference was provided by Natural England

Highlighted reference provided geo-referenced data for a rMCZ or rRA site

Balanced Seas

Abd Aziz, S. A. & Nedwell, D. B. 1986, "The nitrogen cycle of an East Coast, U.K. Saltmarsh: II. Nitrogen fixation, nitrification, denitrification, tidal exchange", *Estuarine, Coastal and Shelf Science*, vol. 22, no. 6, pp. 689-704.

Allison, S. 2012. Assessment of year class and stock levels of European Flat Oyster *Ostrea edulis* in the Ray Sand Channel, part of the Blackwater rMCZ complex.

Anand, P. L. 1937, "An ecological study of the algae of the British chalk-cliffs. Part I", *Journal of Ecology*, vol. 25, no. 1, pp. 153-188.

Anand, P. L. 1937, "A taxonomic study of the algae of the British chalk-cliffs", *Journal of Botany*, vol. Supplement II, 1-51.

Anand, P. L. 1937, "An ecological study of the algae of the British chalk-cliffs. Part II", *Journal of Ecology*, vol. 25, pp. 344-367.

Andrews, C. & Williams, R. B. G. 2000, "Limpet erosion of chalk shore platforms in southeast England", *Earth Surface Processes and Landforms*, vol. 25, no. 12, pp. 1371-1381.

Andrews, C. E. 2001, The measurement of the erosion of the chalk shore platform of East Sussex, the effect of coastal defence structures and the efficacy of macro scale bioerosive agents, in particular the common limpet, *Patella vulgata*, Thesis edn, Sussex University, Brighton AB - Littoral Chalk Communities.

Andrews, M. J. & Wheeler, A. 1985, "Rare And Little-Known Fishes In The Thames Estuary", *Journal of Fish Biology*, vol. 27, no. 1, pp. 59-71.

Anon. 1997, "Oyster boost in Thames", *Fish Farming International*, vol. 24, no. 9, p. 33.

Araujo, F. G., Bailey, R. G., & Williams, W. P. 1998, "Seasonal and between-year variations of fish populations in the middle Thames estuary: 1980-1989", *Fisheries Management and Ecology*, vol. 5, no. 1, pp. 1-21.

Araujo, F. G., Williams, W. P., & Bailey, R. G. 2000, "Fish assemblages as indicators of water quality in the Middle Thames Estuary, England (1980-1989)", *Estuaries*, vol. 23, no. 3, pp. 305-317.

Attrill, M. J. & Rundle, S. D. 2002, "Ecotone or Ecocline: Ecological Boundaries in Estuaries", *Estuarine, Coastal and Shelf Science*, vol. 55, no. 6, pp. 929-936.

Attrill, M. J. & Power, M. 2004, "Partitioning of temperature resources amongst an estuarine fish assemblage", *Estuarine, Coastal and Shelf Science*, vol. 61, no. 4, pp. 725-738.

Bark, A., Williams, B., & Knights, B. 2007, "Current status and temporal trends in stocks of European eel in England and Wales", *ICES Journal of Marine Science*, vol. 64, no. 7, pp. 1368-1378.

Ben Amor, H. 2006, The impact of recreational boat anchoring on the *Zostera marina* in Studland Bay, Bournemouth University.

Bines, T. 2000, North east Kent European marine sites comprising: Thanet coast candidate special area of conservation (cSAC), Thanet coast and sandwich bay special protection area (SPA), sandwich bay candidate special area of conservation (cask), English Nature.

Birchenough, S. N. R., Boyd, S. n. E., Vanstaen, K., Coggan, R. A., & Limpenny, D. S. 2010, "Mapping an aggregate extraction site off the Eastern English Channel: A methodology in support of monitoring and management", *Estuarine, Coastal and Shelf Science*, vol. 87, no. 3, pp. 420-430.

Cefas 2001, the impact of disposal of marine dredged material on the Thanet coast and sandwich bay candidate special areas of conservation (cSACs), CEFAS, Lowestoft (UK).

Coggan, R., Diesing, M., & K., V. 2009, mapping annex i reefs in the central English channel: evidence to support the selection of candidate sacs, Scientific Series Technical Report, Cefas, Lowestoft.

Colclough, S., Fonseca, L., Astley, T., Thomas, K., & Watts, W. 2005, "Fish utilisation of managed realignments", *Fisheries Management and Ecology*, vol. 12, no. 6, pp. 351-360.

Cole, H. A. 1942, "The American whelk tingle, *Urosalpinx cinerea* (Say), on British oyster beds", *Journal of the Marine Biological Association of the United Kingdom*, vol. 25, no. 3, pp. 477-508.

Collins, K. & Mallinson, J. 1988, Marine flora and fauna off Bembridge, Isle of Wight: Sublittoral survey and review of existing knowledge Nature Conservancy Council, Peterborough.

Collins, K. & Mallinson, J. 1988, "Echiurid (*Maxmuelleria lankasteri* (Herdman, 1898)) population off Bembridge, IOW", *Porcupine Newsletter*, vol. 4, p. 56.

Collins, K. 2003, Dorset marine habitat surveys: maerl, worm reefs, bream nests, sea fans and brittlestars. 2003 survey results, Report to Dorset Wildlife Trust and English Nature edn, Dorset Wildlife Trust.

Collins, K. 2003, Dorset marine habitat surveys: maerl, worm reefs, bream nests, sea fans and brittlestars. 2002/ 2003 survey results, Dorset Wildlife Trust, Report to Dorset Wildlife Trust and English Nature.

Collins, K. & Mallinson, J. 2004, "Dorset maerl", Newsletter. Porcupine Marine Natural History Society, vol. 16, pp. 27-31.

Collins, K. 2008, Poole harbour channel post-deepening EIA studies 2006-8. report to Poole harbour commissioners, National Oceanography Centre.

Coulson, M. G., Budd, J. T. C., Withers, R. G., & Nicholls, D. N. 1980, "The shore environment. Vol.1. Methods Remote sensing and field sampling of mudflat organisms in Langstone and Chichester Harbours, Southern England," J. H. Price, D. E. G. Irvine, & W. F. Farnham, eds., Academic Press, pp. 241-263.

Critchley, A. T. 1983, The establishment and increase of *Sargassum muticum* (Yendo) Fensholt populations within the Solent area of southern Britain I. An investigation of the increase in number of population individuals, 26 edn, Walter de Gruyter, Berlin/New York Berlin, New York.

Davidson, P. 1976, "Oyster fisheries of England and Wales", CEFAS.

Diesing, M., Coggan, R., & Vanstaen, K. 2009, "Widespread rocky reef occurrence in the central English Channel and the implications for predictive habitat mapping", Estuarine, Coastal and Shelf Science, vol. 83, no. 4, pp. 647-658.

Dines, R. A. & Wharfe, J. R. 1985, "The environmental impact of paper mill waste discharges to the Swale", Environmental Pollution. Series A, vol. 38, pp. 245-260.

Downburst, U. & Robinson, D. A. 2011, "Block removal and step back wearing as erosion processes on rock shore platforms: a preliminary case study of the chalk shore platforms of south-east England", Earth Surface Processes and Landforms, vol. 36, no. 5, pp. 661-671.

Downburst, U., Williams, R. B. G., Moses, C. A., & Robinson, D. A. 2008, "Foreshore Narrowing along the Coast of Southeast England, UK-A Re-evaluation", Journal of Coastal Research, vol. 24, no. sp3, pp. 14-24.

Emmerson, M. 2000, "Remedial habitat creation: does *Nereis diversicolor* play a confounding role in the colonisation and establishment of the pioneering saltmarsh plant, *Spartina anglica*?", Helgoland Marine Research, vol. 54, no. 2, pp. 110-116.

Evans, J. 1993, "Temporal and spatial variation in low tide counts of wildfowl and waders on selected estuaries in Great Britain", BTO Research Report, vol. 122, p. 50p.

Foden, J., Rogers, S. I., & Jones, A. P. 2009, "Recovery rates of UK seabed habitats after cessation of aggregate extraction", Marine Ecology Progress Series, vol. 390, pp. 15-26.

Foote, Y., Plessis, E., Robinson, D., Henaff, A., & Costa, S. 2006, "Rates and patterns of downwearing of chalk shore platforms of the Channel: comparisons between France and England", Zeitschrift für Geomorphologie, vol. 144, pp. 93-115.

Francis, R. A., Hoggart, S. P. G., Gurnell, A. M., & Coode, C. 2008, "Meeting the challenges of urban river habitat restoration: developing a methodology for the River Thames through central London", *Area*, vol. 40, no. 4, pp. 435-445.

Garbutt, R. A., Reading, C. J., Wolters, M., Gray, A. J., & Rothery, P. 2006, "Monitoring the development of intertidal habitats on former agricultural land after the managed realignment of coastal defences at Tollesbury, Essex, UK", *Marine Pollution Bulletin*, vol. 53, no. 1–4, pp. 155-164.

George, J. D., Tittley, I., & Price, J. H. 1989, *Macrobenthos of chalk shores in the Isle of Wight, Dorset and Devon*, Unpub. NCC CSD Rep. 931 edn, Nature Conservancy Council, Peterborough AB - Littoral Chalk Communities.

Green, B., Smith, D., Grey, J., & Underwood, G. 2012, "High site fidelity and low site connectivity in temperate salt marsh fish populations: a stable isotope approach", *Oecologia*, vol. 168, no. 1, pp. 245-255.

Green, B. C., Smith, D. J., Earley, S. E., Hepburn, L. J., & Underwood, G. J. C. 2009, "Seasonal changes in community composition and trophic structure of fish populations of five salt marshes along the Essex coastline, United Kingdom", *Estuarine, Coastal and Shelf Science*, vol. 85, no. 2, pp. 247-256.

Green, M. & Alderman, D. J. 1983, "Neoplasia in *Mytilus edulis* L. from United Kingdom waters", *Aquaculture*, vol. 30, no. 1-4, pp. 1-10.

Greensmith, J. T. & Tucker, E. V. 1966, "Morphology and evolution of inshore shell ridges and mud mounds in modern intertidal flats, near Bradwell, Essex", *Proceedings of the Geologists Association*, vol. 77, pp. 329-346.

Hassan, H. M. 1994, *Effects of salinity on the plant species of chalk sea-cliffs near Folkstone, Kent*, London University.

Havard, M. S. C. & Tindal, E. C. 1994, "The impacts of bait digging on the polychaete fauna of the Swale Estuary, Kent, UK", *Polychaete Research*, vol. 16, pp. 32-36.

Herbert, R. J. H., Farnham, W., Farnham, T., & Luxton, R. 2012, *status and sensitivity of the bap priority marine alga *padina pavonica*. final report*, SITA TRUST. Enriching Nature. Medina Valley Centre for Environmental & Outdoor Education.

Hill, T. O., Emblow, C. S., & Northen, K. O. 1996, *Marine nature conservation review. Sector 6. Inlets in eastern England: area summaries*, *Coasts and Seas of the United Kingdom MNCR series* edn, Joint Nature Conservation Committee, Peterborough.

Hughes, R. G., Lloyd, D., Ball, L., & Emson, D. 2000, "The effects of the polychaete *Nereis diversicolor* on the distribution and transplanting success of *Zostera noltii*", *Helgoland Marine Research*, vol. 54, no. 2, pp. 129-136.

Hutchinson, P. 1983, "A note recording the occurrence of hermaphroditic smelt, *Osmerus eperlanus* (L.) from River Thames, England", *Journal of Fish Biology*, vol. 23, pp. 241-243.

Irving, RA. 1999. Report of the Sussex SEASE/IRCH Project, 1992-1998. Published by the Sussex SEASEARCH Project. English Nature, Lewes.

James, J. W. C., Coggan, R. A., Blyth-Skyrme, V. J., Morando, A., Birchenough, S. N. R., Bee, E., Limpenny, D. S., Verling, E., Vanstaen, K., Pearce, B., Johnston, C. M., Rocks, K. F., Philpott, S. L., & Rees, H. L. 2007, eastern English channel marine habitat map, Science Series Technical Report, Cefas, Lowestoft.

Jones, L. A. 1999, *Studies of littoral algae of the Isle of Wight and Solent region*, Portsmouth University.

Kamenos, N. A., Moore, P. G., & Hall-Spencer, J. M. 2004, "Nursery-area function of maerl grounds for juvenile queen scallops *Aequipecten opercularis* and other invertebrates", *Marine Ecology Progress Series*, vol. 274, pp. 183-189.

Kenny, A. J. & Rees, H. L. 1994, "The effects of marine gravel extraction on the macrobenthos: Early post-dredging recolonization", *Marine Pollution Bulletin*, vol. 28, no. 7, pp. 442-447.

Kenny, A. J. & Rees, H. L. 1996, "The effects of marine gravel extraction on the macrobenthos: Results 2 years post-dredging", *Marine Pollution Bulletin*, vol. 32, no. 8, pp. 615-622.

Lee, S. & Whitfield, P. J. 1992, "Virus-associated spawning papillomatosis in smelt, *Osmerus eperlanus* L., in the River Thames", *Journal of Fish Biology*, vol. 40, no. 4, pp. 503-510.

Lees, R. G., Rees, H. L., Lambert, M. A., Rowlatt, S. M., & Limpenny, D. S. 1990, "Benthic studies in relation to dredging activity off the Isle of Wight, Southern England", ICES, Copenhagen (Denmark). 1990.

Livingstone, D.R.; Mitchelmore, C.L.; Peters, L.D.; O'Hara, S.C.M.; Shaw, J.P.; Chesman, B.S.; Doyotte, A.; McEvoy, J.; Ronisz, D.; Larsson, D.G.J.; Förlin, L. 2000 "Development of hepatic CYP1A and blood vitellogenin in eel (*Anguilla anguilla*) for use as biomarkers in the Thames Estuary UK" *Marine Environmental Research* vol. 50 issue 1-5 July, 2000. p. 367-371

Mallinson, J. 2001, "The Porcupine Recording Scheme", *Newsletter of the Porcupine Marine Natural History Society*, vol. 7, pp. 12-14.

Möller, I. 2006, "Quantifying saltmarsh vegetation and its effect on wave height dissipation: Results from a UK East coast saltmarsh", *Estuarine, Coastal and Shelf Science*, Volume. 69. Issues 3-4, September 2006, Pages 337-351

McDonald, K. 1985, "A whiter shade of pale", *Diver*, vol. 30, no. 8, pp. 38-39.

Milsom, T. P., Hart, J. D., Parkin, W. K., & Peel, S. 2002, "Management of coastal grazing marshes for breeding waders: the importance of surface topography and wetness", *Biological Conservation*, vol. 103, no. 2, pp. 199-207.

Naismith, I. A. & Knights, B. 1990, "Modelling of unexploited and exploited populations of eels, *Anguilla anguilla* (L.), in the Thames Estuary", *Journal of Fish Biology*, vol. 37, no. 6, pp. 975-986.

Naismith, I. A. & Knights, B. 1993, "The distribution, density and growth of the European eel, *Anguilla anguilla*, in the freshwater catchment of the River Thames", *Journal of Fish Biology*, vol. 42, no. 2, pp. 217-226.

Norton, J., Rollinson, D., & Lewis, J. W. 2005, "Epidemiology of *Anguillicola crassus* in the European eel (*Anguilla anguilla*) from two rivers in southern England", *Parasitology*, vol. 130, no. 06, pp. 679-686.

Paramor, O. A. L. & Hughes, R. G. 2004, "The effects of bioturbation and herbivory by the polychaete *Nereis diversicolor* on loss of saltmarsh in south-east England", *Journal of Applied Ecology*, vol. 41, no. 3, pp. 449-463.

Parker-Humphreys, M. 2005, distribution and relative abundance of demersal fishes from beam trawl surveys in the eastern English channel (ICES division viid) and the southern north sea (ICES division ivc) 1993-2001.

Paul, M. & Amos, C. L. 2011, "Spatial and seasonal variation in wave attenuation over *Zostera noltii*", *Journal of Geophysical Research-Oceans*, vol. 116.

Peters, L. D., Doyotte, A., Mitchelmore, C. L., McEvoy, J., & Livingstone, D. R. 2001, "Seasonal variation and estradiol-dependent elevation of Thames estuary eel *Anguilla anguilla* plasma vitellogenin levels and comparisons with other United Kingdom estuaries", *Science of the Total Environment*, vol. 279, no. 1-3, pp. 137-150.

Pickett, G. D., Brown, M., Harley, B., & Dunn, M. R. 2002, surveying fish populations in the Solent and adjacent harbours using the Cefas bass trawl, CEFAS, Lowestoft.

Pilcher, M. W. & Moore, J. F. 1993, "Distribution and prevalence of *Anguillicola crassus* in eels from the tidal Thames catchment", *Journal of Fish Biology*, vol. 43, no. 3, pp. 339-344.

Pinn, E. H., Richardson, C. A., Thompson, R. C., & Hawkins, S. J. 2005, "Burrow morphology, biometry, age and growth of piddocks (Mollusca : Bivalvia : Pholadidae) on the south coast of England", *Marine Biology*, vol. 147, no. 4, pp. 943-953.

Pinn, E. H., Thompson, R. C., & Hawkins, S. J. 2008, "Piddocks (Mollusca : Bivalvia : Pholadidae) increase topographical complexity and species diversity in the intertidal", *Marine Ecology-Progress Series*, vol. 355, pp. 173-182.

Pinnegar, J. K., Stelzenmüller V., Van Der Kooij, J., Engelhard, G. H., Garrick-Maidment, N., & Righton, D. A. 2008, "Occurrence of the short-snouted seahorse *Hippocampus hippocampus* in the central North Sea", *Cybium*, vol. 32, no. 4, pp. 343-346.

Power, M. & Attrill, M. J. 2007, "Temperature-dependent temporal variation in the size and growth of Thames estuary smelt *Osmerus eperlanus*", *Marine Ecology Progress Series*, vol. 330, pp. 213-222.

Price, J. H. & Tittley, I. 1972, "The marine flora of the county of Kent, southeast England, and its distribution, 1597-1970", *International Seaweed Symposia*, vol. 7, (Proceedings), pp. 31-34.

Rees, H. L., Waldock, R., Matthiesen, P., & Pendle, M. A. 1998, Improvements in the fauna of the Crouch estuary (United Kingdom) following a decline in TBT concentrations ICES, Copenhagen (Denmark).

Richardson, C. A., Collis, S. A., Ekaratne, K., Dare, P., & Key, D. 1993, "The age determination and growth rate of the European flat oyster, *Ostrea edulis*, in British waters determined from acetate peels of umbo growth lines", *ICES Journal of Marine Science: Journal du Conseil*, vol. 50, no. 4, pp. 493-500.

Robinson, D. A. & Jerwood, L. C. 1987, "Frost And Salt Weathering Of Chalk Shore Platforms Near Brighton, Sussex, UK", *Transactions of the Institute of British Geographers*, vol. 12, no. 2, pp. 217-226.

Robinson, D. A. & Jerwood, L. C. 1987, "Sub-aerial weathering of chalk shore platforms during harsh winters in Southeast England", *Marine Geology*, vol. 77, no. 1-2, pp. 1-14.

Robson, E. A. 1957, "A sea-anemone from brackish water", *Nature*, vol. 179, pp. 787-788.

Rogers, S. I. & Millner, R. S. 1996, "Factors affecting the annual abundance and regional distribution of English inshore demersal fish populations: 1973 to 1995", *ICES Journal of Marine Science: Journal du Conseil*, vol. 53, no. 6, pp. 1094-1112.

Ruddock, P. J., Bird, D. J., McEvoy, J., & Peters, L. D. 2003, "Bile metabolites of polycyclic aromatic hydrocarbons (PAHs) in European eels *Anguilla anguilla* from United Kingdom estuaries", *Science of the Total Environment*, vol. 301, no. 1-3, pp. 105-117.

Sheader, M. & Sheader, A. 1987, Lagoon survey of East Sussex and Kent, August, 1987. Final report, Unpub. NCC CSD Rep. 811 edn, Nature Conservancy Council, Peterborough.

Sheader, M. & Sheader, A. 1987, A lagoon survey of the Isle of Wight: final report, Unpub. NCC CSD Rep. 820 edn, Nature Conservancy Council, Peterborough AB - *N. vectensis*; *G. insensibilis*.

Stuttard, P. & Boston, R. N. The Swale National Nature Reserve. *Transactions. Kent Field Club*. 181-189. 1978. The Swale National Nature Reserve. Ref Type: Unpublished Work

Tittley, I. & Spurrier, C. J. H. "Biological Survey of the Intertidal Chalk Reefs between Kingsdown, Deal and Folkestone Warren, Kent".

Tittley, I., Spurrier, C.J.H., Ferrero, T.J., Chimonides, P.J. (2010) Biological survey of the intertidal chalk reef at Seaford to Beachy Head and Brighton to Newhaven Cliffs Site of Special Scientific Interest (SSSI) to set a baseline for SSSI condition assessment. Contract No. FST20/75/026

Tittley, I. 1970, "Kent coast in 1969", Marine Pollution Bulletin, vol. 1, no. 6, p. 83.

Tittley, I. 1972, "Wildlife conservation in the north Kent marshes:: Report of a Working Party.(Nature Conservancy, London). xii+ 42+ liii pp., Marine Pollution Bulletin, vol. 3, no. 6, p. 95.

Tittley, I. & Price, J. H. 1978, "The benthic marine algae of the eastern English Channel: a preliminary floristic and ecological account", Botanica marina, vol. 21, no. 8, pp. 499-512.

Tittley, I. 1985, "Seaweed communities on the artificial coastline of south-eastern England. 1. Reclaimed saline wetland and estuaries", Transactions of the Suffolk Naturalists' Society, vol. 21, pp. 54-64.

Tittley, I. Chalk cliff algal communities of Kent and Sussex, southeast England. [Internal report, No. NC 200D]. 1985. Contractor: British Museum (Natural History), Department of Botany, London. Unpublished report to Nature Conservancy Council (England South-east Region). Chalk cliff algal communities of Kent and Sussex, southeast England. Ref Type: Unpublished Work

Tittley, I. 1985, "Zonation and seasonality of estuarine benthic algae: artificial embankments in the River Thames", Botanica marina, vol. 28, no. 1, pp. 1-8.

Tittley, I. 1985, Chalk-cliff algal communities of Kent and Sussex, southeast England, Unpub. NCC CSD Rep. edn, Nature Conservancy Council, Peterborough AB - Littoral Chalk Communities.

Tittley, I. 1986, "Seaweed communities of the artificial coastline of southeastern England 2. Open sea-shore", Transactions of the Kent Field Club, vol. 10, pp. 55-67.

I.Tittley, B. Chapman, B. Hitchin, C.J.H. Spurrier, T.Child "Thanet Coast Special Area of Conservation 2011 Intertidal Monitoring Report". 2012

Tittley, I., Price, J. H., Fincham, A. A., & George, J. D. 1986, The macrobenthos of chalk and greensand shores in southeastern England. (Flora and fauna), Unpublished NCC CSD Report No. 677 edn, British Museum (Natural History), London AB - Littoral Chalk Communities.

Tittley, I., Price, J. H., Fincham, A. A., & George, J. D. 1986, the macrobenthos of chalk and greensand shores in southeastern England, Nature Conservancy Council. Contractor: British Museum (Natural History), London..

Tittley, I. 1988, Chalk cliff algal communities: 2. Outside southeastern England, Unpub. NCC CSD Rep. 878 edn, Nature Conservancy Council, Peterborough AB - Littoral Chalk Communities.

Tittley, I., George, J. D., Evans, N. J., & Chimonides, P. J. 1989, macrobenthos of non-chalk, rocky shores in southeastern England, Contractor: British Museum (Natural History, London.) Nature Conservancy Council.

Tittley, I., George, J. D., Evans, N. J., & Chimonides, P. J. 1989, Macrobenthos of non-chalk rocky shores in southeastern England, Unpub. NCC CSD Rep. 1005 edn, Nature Conservancy Council, Peterborough AB - Littoral Chalk Communities.

Tittley, I., Gilliland, P., & Pound, D. 1998, "Marine flora of the Thanet coast marine SAC: A conservation and management perspective," in Changes in the marine flora of the North Sea, G. W. Scott & I. Tittley, eds., Centre for Environmental Research into Coastal Issues (CERCI), Scarborough, pp. 65-75.

Tittley, I. 2002, "Marine algal (seaweed) collections at the Natural History Museum, London (BM): past, present and future", Constancea: University of California Electronic Publications in Botany, vol. 83.

Tittley, I. 1971, "The Kent coast in 1970", Marine Pollution Bulletin, vol. 2, no. 8, pp. 120-122.

Tittley, I. 1972, "The Kent coast in 1971", Marine Pollution Bulletin, vol. 3, no. 9, pp. 135-138.

Tubbs, C. R. & Tubbs, J. M. 1983, "The distribution of *Zostera* and its exploitation by wildfowl in the Solent, Southern England", Aquatic Botany, vol. 15, no. 3, pp. 223-239.

Unicomarine, 2005. Review of data in Stour and Orwell Estuaries HHA report 58

van den Broek, W. L. F. 1979, "A seasonal survey of fish populations in the Lower Medway Estuary, Kent, based on power station screen samples", Estuarine and Coastal Marine Science, vol. 9, no. 1, pp. 1-15.

Whomersley, P., Huxham, M., Schratzberger, M., & Bolam, S. 2009, "Differential response of meio- and macrofauna to in situ burial", Journal of the Marine Biological Association of the United Kingdom, vol. 89, no. 06, pp. 1091-1098.

Williams, C. & Clark, R. 2010, report on the chalk reefs of Sussex, exemplified by the recreational dive sites: south west rocks (msnci), looe gate (msnci) and ship rock (msnci), Natural England.

Williams, C. & Clark, R. C., Jr. 2010, report on the chalk reefs of Sussex, exemplified by the recreational dive sites: south west rocks (msnci), looe gate (msnci) and ship rock (msnci), Report for Sussex Seasearch.

Williams, R. B. 1976, "Conservation of the sea anemone *Nematostella vectensis* in Norfolk, England and its world distribution", Transactions of the Norfolk and Norwich Naturalists Society, vol. 23, pp. 257-266.

Wilson, P. 1999, "*Tenellia adspersa* (Nordmann, 1845) living at St. Osyth in Essex", Journal of Conchology, vol. 36, no. 5, p. 125.

Worsfold, T.M., & Dyer, M.f., 2004. The distribution of birds of ross (*Sabellaria spinulosa*) and other biotopes in Harwich Harbour. Unicomarine Report HHASab04 to Harwich Haven Authority, September 2004

Wyer, D. W., Boorman, L. A., & Waters, R. 1977, "Studies on the distribution of *Zostera* in the outer Thames Estuary", *Aquaculture*, vol. 12, pp. 215-227.

Finding Sanctuary

Amaral, V., Cabral, H. N., Jenkins, S., Hawkins, S., & Paula, J. 2009, "Comparing quality of estuarine and nearshore intertidal habitats for *Carcinus maenas*", Estuarine, Coastal and Shelf Science, vol. 83, no. 2, pp. 219-226.

Bale, A. J., Uncles, R. J., Villena-Lincoln, A., & Widdows, J. 2007, "An assessment of the potential impact of dredging activity on the Tamar Estuary over the last century: Bathymetric and hydrodynamic changes", Hydrobiologia, vol. 588, pp. 83-95.

Bayne, B. L., Hawkins, A. J. S., & Navarro, E. 1987, "Feeding and digestion by the mussel *Mytilus edulis* L. (*Bivalvia: Mollusca*) in mixtures of silt and algal cells at low concentrations", Journal of Experimental Marine Biology and Ecology, vol. 111, no. 1, pp. 1-22.

Belderson, R. H., Wilson, J. B., & Holme, N. A. 1988, "Direct observation of longitudinal furrows in gravel and their transition with sand ribbons of strongly tidal seas", Tide-Influenced Sedimentary Environments and Facies pp. 79-90.

Bird, E. C. F. 1972, "The physiography of the Fleet", Proceedings. Dorset Natural History and Archaeological Society, vol. 93, pp. 125-131.

Bland, S., Ackroyd, Marsh, J. G., & Millward, G. E. 1982, "Heavy Metal Content of Oysters From the Lynher Estuary, U.K.", Science of the Total Environment, vol. 22, no. 3, pp. 235-241.

Blunden, G., Farnham, W. F., Jephson, N., Barwell, C. J., Fenn, R. H., & Plunkett, B. A. 1981, "Proceedings of the Xth International Seaweed Symposium, Goteborg, 11-15 August 1980 The composition of maerl beds of economic interest in northern Brittany, Cornwall and Ireland," T. Levring, ed., Walter de Gruyter, Berlin AB - Maerl; *L. corallioides*; *P. calcareum*, pp. 651-656.

Bosence, D. & Wilson, J. 2003, "Maerl growth, carbonate production rates and accumulation rates in the northeast Atlantic", Aquatic Conservation-Marine and Freshwater Ecosystems, vol. 13, p. S21-S31.

Brown, J. H. & Hunnam, P. J. 1976, "The marine fauna of Lundy Opisthobranchia", Report of Lundy Field Society, vol. 27, pp. 37-47.

Carter, M. C., Bishop, J. D. D., Evans, N. J., & Wood, C. A. 2010, "Environmental influences on the formation and germination of hibernacula in the brackish-water bryozoan *Victorella pavida* Saville Kent, 1870 (*Ctenostomata: Victorellidae*)", Journal of Experimental Marine Biology and Ecology, vol. 383, no. 2, pp. 89-95.

Clark, E. 2011, "Looe Seagrass Mapping report"

Collins, K. & Mallinson, J. 1991, "The Kimmeridge underwater nature trail", Sanctuary, vol. 20, pp. 22-24.

Collins, K., Hall-Spencer, J. M., Grall, J., Moore, P. G., & Atkinson, R. J. A. 2003, "Dorset marine habitat surveys: maerl, worm reefs, bream nests, sea fans and brittlestars. 2003 survey results Bivalve fishing and maerl-bed conservation in France and the UK-retrospect and prospect", Aquatic Conservation: Marine and Freshwater Ecosystems, vol. Report to Dorset Wildlife Trust and English Nature, p. 14p.

Collins, K. 2005, Dorset marine habitat surveys: maerl, worm reefs, brittlestars, sea fans and seagrass. 2004 field report, Progress report to English Nature, Project Ref: DP1/DorsetmarineHabitat/04/06 edn, English Nature.

Cook, K.J. 2011. "Report on 2011 Isles of Scilly *Zostera marina* survey". Report to Natural England.

Corbin, P. G. 1979, "The seasonal abundance of four species of Stauromedusae (*Coelenterata: Scyphomedusae*) at Plymouth", *Journal of the Marine Biological Association of the United Kingdom*, vol. 59, no. 02, pp. 385-391.

Curtis, L. (2010) Littoral biotope survey and Condition Assessment of the Lynher Estuary 2010
Deeble, M. & Stone, V. 1985, "A port that could threaten marine life in England's Fal Estuary", *Oryx*, vol. 19, no. 2, pp. 74-78.

Dixon, I. & Moore, J. 1987, surveys of harbours, rias and estuaries in southern Britain: the Solent system, Report to the Nature Conservancy Council from the Field Studies Council Oil Pollution Research Unit, Orierton Field Centre, Pembroke, Dyfed, UK.

Doyle, J., Lordan, C., Fitzgerald, R., O'Connor, S., Fee, D., Nolan, C., & Hayes, J. 2011, "Celtic Sea Nephrops Grounds 2011 UWTV Survey Report".

Elliott, M. & Mazik, K. 2011, Rame head environmental impact study: review of evidence, Report to Marine Management Organisation. Institute of Estuarine and Coastal Studies University of Hull.

Farnham, W. F. & Jephson, N. A. 1977, "A Survey Of The Maerl Beds Of Falmouth Cornwall England", *British Phycological Journal*, vol. 12, no. 2, p. 119.

Faubel, A. & Warwick, R. M. 2005, "The marine flora and fauna of the Isles of Scilly: Free-living Plathelminthes ('Turbellaria')", *Journal of Natural History*, vol. 39, no. 1, pp. 1-45.

Forster, G. R. 1958, "Underwater observations on the fauna of shallow rocky areas in the neighbourhood of Plymouth", *Journal of the Marine Biological Association of the United Kingdom*, vol. 37, no. 02, pp. 473-482.

Friend, P. L., Velegrakis, A. F., Weatherston, P. D., & Collins, M. B. 2006, "Sediment transport pathways in a dredged ria system, southwest England", *Estuarine, Coastal and Shelf Science*, vol. 67, no. 3, pp. 491-502.

Gall, A. 2011. "Marine BAP Habitats and Species of the Isles of Scilly - an update to the Isles of Scilly Environmental Audit 2008".

Garrick-Maidment, N., Newman, J., & Durant, E. Movement of a pair of Spiny Seahorses (*Hippocampus guttulatus*) seen during the summer of 2010 at Studland Bay in Dorset. 9. 2010. Escot, Devon, The Seahorse Trust. Movement of a pair of Spiny Seahorses (*Hippocampus guttulatus*) seen during the summer of 2010 at Studland Bay in Dorset. Ref Type: Unpublished Work

George, C. L. & Warwick, R. M. 1985, "Annual macrofauna production in a hard-bottom reef community", *Journal of the Marine Biological Association of the United Kingdom*. Plymouth, vol. 65, no. 3, pp. 713-735.

Gill, C. & Mercer, T. 1989, *Surveys of harbours, rias and estuaries in southern Britain: Camel Estuary*. Volume 1, Unpub. NCC CSD Rep. 954 edn, Nature Conservancy Council, Peterborough.

Hall-Spencer, J., Munn, C. B., & Hiscock, K. "Scientific divers quantify first known outbreaks of cold-water coral disease", in *Research in Shallow Marine and Fresh Water Systems*. 1st International Workshop - Proceedings, B. J. Merkel & M. Schipek, eds., Freiberg Online Geology, p. 144.

Hamilton, E. L. 1980, "Geoacoustic modeling of the sea floor", *Journal of the Acoustical Society of America*, vol. 68, no. 5, pp. 1313-1340.

Harvey, L. A. 1969, "Common structures and properties of seagrass beds fringing the coasts of the world", *Journal of Natural History*, vol. 3, no. 1, pp. 3-18.

Harvey, L. A. 1969, "The marine flora and fauna of the Isles of Scilly. The islands and their ecology", *Journal of Natural History*, vol. 3, no. 1, pp. 3-18.

Hawkins, S. J. 1991, "Further reading on the studies of marine life and habitats around Lundy Island".

Herbert, R. J. H., Farnham, W., & Tittley, I. 2010, "Reproduction and habitat of the marine alga *Padina pavonica* (*Dictyotales, Phaeophyceae*) in the British Isles", *Porcupine Newsletter* no. 27, pp. 28-29.

Hiscock, K. 1979, "A survey of sublittoral habitats and species in the region of Padstow, Cornwall", *Progress in underwater Science*. Rep. underwat. Ass.(NS), vol. 4, pp. 121-130.

Hiscock, K. 1994, "Marine communities at Lundy - origins, longevity and change", *Biological Journal of the Linnean Society*, vol. 51, no. 1/2, pp. 183-188.

Holme, N. A. 1950, "The bottom fauna of Great West Bay", *Journal of the Marine Biological Association of the United Kingdom*, vol. 29, pp. 163-183.

Holme, N. A. "Fluctuations in the benthos of the western English Channel".

Holme, N. A. & Wilson, J. B. 1985, "Faunas associated with longitudinal furrows and sand ribbons in a tide-swept area in the English channel", *Journal of the Marine Biological Association of the United Kingdom*, Plymouth, vol. 65, no. 4, pp. 1051-1072.

Horsman, P. 1986, a report on the river dart survey, Nature Conservancy Council (NCC), London (United Kingdom).

Hoskin, M. G., Coleman, R. A., & von Carlshausen, L. 2009, ecological effects of the lundy no-take zone: the first five years (2003-2007). final report to natural England, Defra & Wwf-Uk.

Huvenne, V. A. I., Tyler, P. A., Masson, D. G., Fisher, E. H., Hauton, C., Hühnerbach, V., Le Bas, T. P., & Wolff, G. A. 2011, "A Picture on the Wall: Innovative Mapping Reveals Cold-Water Coral Refuge in Submarine Canyon", *PLoS One*, vol. 6, no. 12, p. e28755.

Irvine, D. E. G., Smith, R. M., Tittley, I., Fletcher, R. L., & Farnham, W. F. 1972, "A survey of the marine algae of Lundy", *British Phycological Journal*, vol. 7, pp. 119-135.

Irving, R. 2004, "Leptopsammia pruvoti at Lundy - teetering on the brink?", *Newsletter. Porcupine Marine Natural History Society*, vol. 15, pp. 29-34.

Irving, R. A. & Hiscock, K. 2010, "The status of the sunset cup coral *Leptopsammia pruvoti* at Lundy", *Journal of the Lundy Field Society*, vol. 2, pp. 67-84.

Irving, R.A. and Northen, K.O. 2012. "Isles of Scilly SAC Diving Monitoring Studies, 2011". Natural England Commissioned Reports, Number 104.

Jackson, E.L., Higgs, S., Allsop, T., Cawthray, A., Evans, J. and Langmead, O. 2011 "Isles of Scilly Seagrass Mapping". Natural England Commissioned Reports, Number 087.

Jones, G. B., Matharu, H. S., & Jones, R. E. H. 1973, "Heavy metals in organisms and sediments from the east coast of Lundy", *Report of the Lundy Field Society* no. 23, pp. 36-39.

Langston, W., Chesman, B., Burt, G., Taylor, M., Covey, R., Cunningham, N., Jonas, P., & Hawkins, S. 2006, "Characterisation of the European Marine Sites in South West England: the Fal and Helford candidate Special Area of Conservation (cask)", *Marine Biodiversity* pp. 321-333.

Light, J. 1986, "Paludinella littorina living along the Fleet, Dorset", *Journal of Conchology*, vol. 32, p. 260.

Lorenz, F. & Melaun, C. 2011, "A new species of Simnia from England (*Caenogastropoda: Ovulidae*)", *Molluscan Research*, vol. 31, no. 3, pp. 167-175.

Lowe, D. M. & Moore, M. N. 1979, "The cytochemical distributions of zinc (Zn II) and iron (Fe III) in the common mussel, *Mytilus edulis*, and their relationship with lysosomes", *Journal of the Marine Biological Association of the United Kingdom*, vol. 59, no. 04, pp. 851-858.

Maggs, C. A. 1990, "Distribution and evolution of non-coralline crustose red algae in the North Atlantic", *Evolutionary biogeography of the marine algae of the North Atlantic* pp. 241-264.

MahÃ©, J. C. & Poulard, J. C. 2005, spatial distribution and abundance trends of main elasmobranch species in the bay of biscay and the celtic sea from bottom trawl surveys.

Mangi, S. C., Rodwell, L. D., & Hattam, C. 2011, "Assessing the Impacts of Establishing MPAs on Fishermen and Fish Merchants: The Case of Lyme Bay, UK", *AMBIO: A Journal of the Human Environment* pp. 1-12.

Mercer, T., Howson, C., & Bunker, F. 2006, Lundy european marine site sublittoral monitoring report 2003/4, Contractor: Aquatic Survey & Monitoring Ltd., Peterborough.

Moore, J., Smith, J., Northen, K., & Little, M. 1998, Inlets in the Bristol Channel and approaches: area summaries. *Marine Nature Conservation Review Sector 9, Coasts and seas of the United Kingdom - MNCR series edn*, Joint Nature Conservation Committee, Peterborough.

Moore, J. J., Smith, J., & Northen, K. O. 1999, *Marine Nature Conservation Review. Sector 8. Inlets in the western English Channel. Area summaries* Joint Nature Conservation Committee.

Mortensen, P. B. & Fosså, J. H. "Species diversity and spatial distribution of invertebrates on deep-water *Lophelia* reefs in Norway", in *Proceedings of 10th International Coral Reef Symposium*, 1849-1868 (2006).

Pirie, B. J. S., George, S. G., Lytton, D. G., & Thomson, J. D. 1984, "Metal-containing blood cells of oysters: ultrastructure, histochemistry and X-ray microanalysis", *Journal of the Marine Biological Association of the United Kingdom*, vol. 64, no. 01, pp. 115-123.

Potts, G. W. & Swaby, S. E. 1995, Species recovery programme. The giant goby (*Gobius cobitis*) in the south west of England *English Nature*, Peterborough AB - *G. cobitis*.

Probert, R. J. & Brenchley, J. L. 1999, "The effect of environmental factors on field and laboratory germination in a population of *Zostera marina* L. from southern England", *Seed Science Research*, vol. 9, no. 04, pp. 331-339.

Robins, M. W. 1969, "The marine flora and fauna of the Isles of Scilly Cnidaria and Ctenophora", *Journal of Natural History*, vol. 3, no. 3, pp. 329-343.

ROYAL HASKONING, 2008. Site Selection Report for the Inshore Marine SACs Project. Salcombe to Yealm & Eddystone Site Selection. Report No. 9SO282/SSR/Salcombe/01

Russell, G. 1968, "List of marine algae from the Isles of Scilly", *British Phycological Bulletin*, vol. 3, no. 3, pp. 579-584.

Schmid, R., Dring, M. J., & Forster, R. M. 1994, "Kinetics Of Blue-Light Stimulation And Circadian Rhythmicity Of Light-Saturated Photosynthesis In Brown Algae: A Species Comparison", *Journal of Phycology*, vol. 30, no. 4, pp. 612-621.

Scourse, J. D., Austin, W. E. N., Bateman, R. M., Catt, J. A., Evans, C. D. R., Robinson, J. E., & Young, J. R. 1990, "Sedimentology and micropalaeontology of glacial marine sediments from the Central and Southwestern Celtic Sea", Geological Society, London, Special Publications, vol. 53, no. 1, pp. 329-347.

Seasearch. 2010, "Isles of Scilly 2010 Summary Report".

Seaward, D. R. 1978, "The marine molluscs of the Fleet, Dorset", Proceedings of the Dorset Natural History and Archaeological Society, vol. 100, pp. 100-108.

Seaward, D. R. 1987, "Caecum armoricum De Folin, 1869, new to the British marine molluscan fauna, living in the Fleet, Dorset, within an unusual habitat", Proceedings Dorset Natural History and Archaeological Society, vol. 109, p. 165.

Seaward, D. R. 1987, "The marine molluscs of Dorset", Proceedings Dorset Natural History and Archaeological Society, vol. 109, p. 164.

Shedder, M. & Shedder, A. 1987, Lagoon survey of Avon and Gloucestershire (Weston-super-Mare to Gloucester), July, 1987. Final report, Unpub. NCC CSD Rep. 813 edn, Nature Conservancy Council, Peterborough.

Shedder, M. & Shedder, A. L. 1987, "The distribution of the lagoonal amphipod *Gammarus insensibilis* Stock in England", Porcupine Newsletter, vol. 3, pp. 220-223.

Smith, P. & Nunny, R. 2012, "Mapping of sedimentary marine biotopes around Lundy, UK", Journal of the Lundy Field Society, vol. 3, pp. 41-74.

Smith, R. I. 1956, "The ecology of the Tamar Estuary. VII Observations on the interstitial salinity of intertidal muds in the estuarine habitat of *Nereis diversicolor*", Journal of the Marine Biological Association of the United Kingdom, vol. 35, pp. 81-104.

Spooner, G. M. 1928, "Fauna of Weymouth Bay as indicated by jetsam on Weymouth beach", Proceedings Dorset Natural History and Archaeological Society, vol. 49, pp. 200-201.

Spooner, G. M. & Moore, H. B. 1940, "The ecology of the Tamar Estuary VI. An account of the macrofauna of the intertidal muds", Journal of the Marine Biological Association of the United Kingdom, vol. 24, no. 01, pp. 283-330.

Sykes, E. R. 1945, "The marine molluscs of Dorset", Proceedings of the Dorset Natural History and Antiquarian Field Club, vol. 67, pp. 127-142.

UNIVERSITY OF PLYMOUTH, 2011. "South Devon reef video baseline surveys for the Prawle Point to Plymouth Sound & Eddystone cSAC and surrounding areas". May, 2011

Vattakaven, T., Bond, P., Bradley, G., & Munn, C. B. 2006, "Differential effects of temperature and starvation on induction of the viable-but-nonculturable state in the coral pathogens *Vibrio shiloi* and *Vibrio tasmaniensis*", *Applied and environmental microbiology*, vol. 72, no. 10, pp. 6508-6513.

Warne, S. & Jones, B. W. 1995, species distributions from the English celtic sea groundfish surveys, 1984 to 1991.

Wheeler, A. W. 1960, "Gobius capito as a British fish", *Annals and magazine of Natural History, Series 13*, vol. 3, no. 27, pp. 177-181.

Wilson, D. P. 1970, "Additional observations on larval growth and settlement of *Sabellaria alveolata*", *Journal of the Marine Biological Association of the United Kingdom*, vol. 50, no. 1, pp. 1-31.

Woods, C. 2008, "Seasearch pink sea fan surveys 2004/6", A report for the Marine Conservation Society.

Irish Sea

Allen JH, Billings I, Cutts N & Elliott M. 2002. Mapping, condition and conservation assessment of honeycomb worm *Sabellaria alveolata* reefs on the Eastern Irish Sea coast. Report to English Nature. Institute of Estuarine and Coastal Studies, University of Hull. November 2002. Report no. Z122-F-2002

Allen JH & Hemingway KL. 2005. The Dee Estuary biotope survey 2004/5. Report to English Nature. Institute of Coastal and Estuarine Studies, Hull. Report no. ZBB640-F-2005

Allen, J. R., Wilkinson, S. B., Hawkins, S. J., Hartnoll, R. G., & Russell, G. A. 1991, baseline survey of the cumbrian shores from st. bees head to maryport, final report.

CMACS. 2011. North West Region European marine sites: Condition monitoring of Littoral Features. Report to Natural England. Centre for Marine and Coastal Studies. Report reference: J3155

Fish, J. & Fish, S. 2011, A student's guide to the seashore, 3, illustrated edn, Cambridge University Press.

Frost, M. T., Leaper, R., Mieszkowska, N., Moschella, P., Murua, J., Smyth, C., & Hawkins, S. J. 2004, "ID 203.pdf", Occasional Publications. Marine Biological Association of the United Kingdom, vol. 16, p. 57p.

Frost, M. T., Leaper, R., Mieszkowska, N., Moschella, P., Murua, J., Smyth, C., & Hawkins, S. J. 2004, Recovery of a biodiversity action plan species in northwest England: possible role of climate change, artificial habitat and water quality amelioration MBA, Plymouth (UK).

Hill, A. S., Brand, A. R., Veale, L. O., & Hawkins, S. J. 1997, "Assessment of the effects of scallop dredging on benthic communities", Contract CSA, vol. 2332.

Hill, A. S., Veale, L. O., Pennington, D., Whyte, S. G., Brand, A. R., & Hartnoll, R. G. 1999, "Changes in Irish Sea benthos: possible effects of 40 years of dredging", *Estuarine, Coastal and Shelf Science*, vol. 48, no. 6, pp. 739-750.

Kaiser, M. et al (2002) Predicting the displacement of Common Scoter (*Melanitta nigra*) from benthic feeding areas due to offshore wind farms. Centre for Applied Marine Sciences, UCNW, Bangor

Mackie, A. S., Oliver, P. G., & Rees, E. I. S. 1995, benthic biodiversity in the southern Irish sea.

Mills, D. J. L. 1998, "Cardigan Bay and north Wales (Cwm-yr-Eglwys, Newport Bay to Rhŷs-on-Sea) (MNCR Sector 10)," in *Marine Nature Conservation Review. Benthic marine ecosystems of Great Britain and the north-east Atlantic, Coasts and seas of the United Kingdom*. MNCR series edn, K. Hiscock, ed., Joint Nature Conservation Committee, Peterborough, pp. 297-314.

Parker-Humphreys, M. 2004, distribution and relative abundance of demersal fishes from beam trawl surveys in the Irish sea (ICES Division viia) 1993-2001, CEFAS, Lowestoft.

Robinson, K. A., Mackie, A. S. Y., Lindenbaum, C., Darbyshire, T., van Landeghem, K. J. J., & Sanderson, W. G. 2012, "37 - Seabed Habitats of the Southern Irish Sea," in Seafloor Geomorphology as Benthic Habitat, Elsevier, London AB - Modiolus Modiolus Beds, pp. 523-537.

Skelcher, G. & Warton, C. 2010, Ribble estuary nnr management plan.

Veale, L. O., Hill, A. S., Hawkins, S. J., & Brand, A. R. 2000, "Effects of long-term physical disturbance by commercial scallop fishing on subtidal epifaunal assemblages and habitats", *Marine Biology*, vol. 137, no. 2, pp. 325-337.

Veale, L. O., Hill, A. S., Hawkins, S. J., & Brand, A. R. 2001, "Distribution and damage to the by-catch assemblages of the northern Irish Sea scallop dredge fisheries", *Journal of the Marine Biological Association of the United Kingdom*, vol. 81, no. 1, pp. 85-96.

Net Gain

Austen, M. C., Buchanan, J. B., Hunt, H. G., Josepson, A. B., & Kendall, M. A. 1991, "Comparison of long-term trends in benthic and pelagic communities of the North Sea", *Journal of the Marine Biological Association of the United Kingdom*, vol. 71, no. 1, pp. 179-190.

Basford, D. J., Eleftheriou, A., Davies, I. M., Irion, G., & Soltwedel, T. 1993, "The ICES North Sea benthos survey: the sedimentary environment", *ICES Journal of Marine Science: Journal du Conseil*, vol. 50, no. 1, pp. 71-80.

Begum, S., Basova, L., Strahl, J., Sukhotin, A., Heilmayer, O., Philipp, E., Brey, T., & Abele, D. 2009, "A Metabolic Model for the Ocean Quahog *Arctica islandica*. "Effects of Animal Mass and Age, Temperature, Salinity, and Geography on Respiration Rate", *Journal of Shellfish Research*, vol. 28, no. 3, pp. 533-539.

Bennett, T. L. & Foster-Smith, J. L. 1998, "South-east Scotland and north-east England (Dunbar to Bridlington) (MNCR Sector 5)," in *Marine Nature Conservation Review. Benthic marine ecosystems of Great Britain and the north-east Atlantic, Coasts and seas of the United Kingdom*. MNCR series edn, K. Hiscock, ed., Joint Nature Conservation Committee, Peterborough, pp. 155-177.

Bucke, D. 1994, *Monitoring for diseases in marine and freshwater fish*, 1991.

Centrica, S. p. *Rough Offshore Facilities Environmental Statement 2008*. 14. 2009. Centrica Storage plc. *Rough Offshore Facilities Environmental Statement 2008*. Ref Type: Unpublished Work

Collie, J. S., Hall, S. J., Kaiser, M. J., & Poiner, I. R. 2000, "A quantitative analysis of fishing impacts on shelf-sea benthos", *Journal of Animal Ecology*, vol. 69, no. 5, pp. 785-798.

Daan, N. 2006, spatial and temporal trends in species richness and abundance for the southerly and northerly components of the north sea fish community separately, based on ibts data 1977-2005 500.

Dyer, M. F., Fry, W. G., Fry, P. D., & Cranmer, G. J. 1982, "A series of North Sea benthos surveys with trawl and headline camera", *Journal of the Marine Biological Association of the United Kingdom*, vol. 62, no. 2, pp. 297-312.

Dyer, M. F., Fry, W. G., Fry, P. D., & Cranmer, G. J. 1983, "Benthic regions within the North Sea", *Journal of the Marine Biological Association of the United Kingdom*, vol. 63, no. 03, pp. 683-693.

English, N. 2000, *Wash and North Norfolk Coast: European Marine Site: English Nature's Advice Given Under Regulation 32 (2) of the Conservation (Nature Habitats &c.) Regulations 1994* English Nature.

Foster-Smith, J. 2000, *The marine fauna and flora of the Cullercoats District. Marine species records for the North East coast of England* University of Newcastle, Department of Marine Sciences and Coastal Management, Newcastle upon Tyne.

Foster-Smith, R. L. 1989, A survey of boulder habitats on the Northumberland coast with a discussion on survey methods for boulder habitats, Unpub. NCC CSD Rep. 921 edn, Nature Conservancy Council, Peterborough.

Frid, C. L. J. 1987, The sub-littoral chalk reefs of North Norfolk: a preliminary survey, Unpub. NCC CSD Rep. 789 edn, Nature Conservancy Council, Peterborough AB - Littoral Chalk Communities.

Frid, C. L. J., Buchanan, J. B., & Garwood, P. R. 1996, "Variability and stability in benthos: twenty-two years of monitoring off Northumberland", ICES Journal of Marine Science: Journal du Conseil, vol. 53, no. 6, pp. 978-980.

Frid, C. L. J., Harwood, K. G., Hall, S. J., & Hall, J. A. 2000, "Long-term changes in the benthic communities on North Sea fishing grounds", ICES Journal of Marine Science: Journal du Conseil, vol. 57, no. 5, pp. 1303-1309.

George, J. D., Tittley, I., Price, J. H., & Fincham, A. A. 1988, The macrobenthos of chalk shores in North Norfolk and around Flamborough Headland (North Humberside), Unpub. NCC CSD Rep. 833 edn, Nature Conservancy Council, Peterborough AB - Littoral Chalk Communities.

George, J. D., Chimonides, P. J., Evans, N. J., & Muir, A. L. "Fluctuations in the macrobenthos of a shallow-water cobble habitat off North Norfolk, England", in Biology and ecology of shallow coastal waters: proceedings of the 28th European Marine Biology Symposium, Institute of Marine Biology of Crete, Iraklio, Crete, 1993, A. Eleftheriou, C. Smith, & A. D. Ansell, eds., Olsen & Olsen, Crete AB - Littoral Chalk Communities, p. 167.

Gilliland, P. M. & Sanderson, W. G. 2000, "Re-evaluation of marine benthic species of nature conservation importance: A new perspective on certain 'lagoonal specialists' with particular emphasis on *Alkmaria romijni* Horst (Polychaeta: Ampharetidae)", Aquatic Conservation: Marine and Freshwater Ecosystems, vol. 10, no. 1, pp. 1-12.

Greenstreet, S. P. R., Spence, F. E., & McMillan, J. A. 1999, "Fishing effects in northeast Atlantic shelf seas: patterns in fishing effort, diversity and community structure. V. Changes in structure of the North Sea groundfish species assemblage between 1925 and 1996", Fisheries Research, vol. 40, no. 2, pp. 153-183.

Hiddink, J. G., Jennings, S., Kaiser, M. J., Queiros, A. M., Duplisea, D. E., & Piet, G. J. 2006, "Cumulative impacts of seabed trawl disturbance on benthic biomass, production, and species richness in different habitats", Canadian Journal of Fisheries and Aquatic Sciences, vol. 63, no. 4, pp. 721-736.

KrÄncke, I. 1995, "Long-term changes in North Sea benthos", Senckenbergiana maritima.Frankfurt/Main, vol. 26, no. 1, pp. 73-80.

Lindeboom, H. J. & De Groot, S. J. 1998, "IMPACT-II: The effects of different types of fisheries on the North Sea and Irish Sea benthic Ecosystems", NIOZ-Rapport, vol. 1998, no. 1.

Moore, P. G. 1973, "The kelp fauna of northeast Britain. I. Introduction and the physical environment", *Journal of Experimental Marine Biology and Ecology*, vol. 13, no. 2, pp. 97-125.

Murby, P. 2008, the wash natural area profile, English Nature.

Rees, H. L. & Nicholson, M. D. 1989, "Trends in lead levels in the horse-mussel from the western North Sea", *Marine Pollution Bulletin*, vol. 20, no. 2, pp. 86-89.

Rees, H. L., Rowlett, S. M., Lambert, M. A., Lees, R. G., & Limpenny, D. S. 1992, "Spatial and temporal trends in the benthos and sediments in relation to sewage sludge disposal off the northeast coast of England", *ICES Journal of Marine Science: Journal du Conseil*, vol. 49, no. 1, pp. 55-64.

Richardson, C. A., Chensery, S. R. N., & Cook, J. M. 2001, "Assessing the history of trace metal (Cu, Zn, Pb) contamination in the North Sea through laser ablation - ICP-MS of horse mussel *Modiolus modiolus* shells", *Marine Ecology Progress Series*, vol. 211, pp. 157-167.

Salmon, K. 2011, york field development project. offshore environmental statement addendum, Centrica Resources Limited, Revision: 02.

Sanchez-Hernandez, C., Boyd, D. S., & Foody, G. M. 2007, "Mapping specific habitats from remotely sensed imagery: Support vector machine and support vector data description based classification of coastal saltmarsh habitats", *Ecological Informatics*, vol. 2, no. 2, pp. 83-88.

Schoene, B. R., Kroencke, I., Houk, S. D., Freyre Castro, A. D., & Oschmann, W. 2003, "The Cornucopia of Chilly Winters: Ocean Quahog (*Arctica islandica* L., Mollusca) master chronology reveals bottom water nutrient enrichment during colder winters (North Sea)", *Senckenbergiana Maritima*, vol. 32, no. 1-2, pp. 165-175.

Schone, B. R., Kroncke, I., Houk, S. D., Castro, A. D. F., & Oschmann, W. 2003, "The Cornucopia of chilly winters: ocean quahog (*Arctica islandica* L., Mollusca) master chronology reveals bottom water nutrient enrichment during colder winters (North Sea)", *Senckenbergiana Maritima*, vol. 32, no. 1/2, pp. 165-175.

Scira Offshore Energy Ltd. Environmental Statement of the Sheringham Shoal wind farm. [SCIRA-5-15-3-MT-RP-02555-A1-ES], 722. 2006. Scira Offshore Energy Ltd. Environmental Statement of the Sheringham Shoal wind farm. Ref Type: Unpublished Work

Spray, R. & Watson, D. Seasearch East, 2010. Marine surveys conducted by Seasearch East. 12. 2011. Seasearch East. Seasearch East, 2010. Marine surveys conducted by Seasearch East. Ref Type: Unpublished Work

Tittley, I. 1998, "Medium to long-term studies on the rocky shore vegetation of North Norfolk," in *Changes in the Marine Flora of the North Sea*, G. W. Scott & I. Tittley, eds., Centre for Environmental Research into Coastal Issues, Scarborough, pp. 53-63.

Vine, H. 1999, "Development of Blue Mussels on the Wash/North Norfolk coast", Waddensea Newsletter, vol. 1, pp. 36-38.

Walker, P., Dare, P. J., & Clark, E. 1990, The potential of aerial photography for estimating intertidal stocks of mussels (*Mytilus edulis*) in the Wash, England ICES, Copenhagen (Denmark).

Walker, P. A. & Heessen, H. J. L. 1996, Long-term changes in ray populations in the North Sea Academic Press, London (UK).

Williams, R. B. 2003, "Locomotory behaviour and functional morphology of *Nematostella vectensis* (Anthozoa: Actiniaria: Edwardsiidae): a contribution to a comparative study of burrowing behaviour in athenarian sea anemones", *Zoologische Verhandelingen*, vol. 345, pp. 437-484.

Witbaard, R., Jenness, M. I., Van Der Borg, K., & Ganssen, G. 1994, "Verification of annual growth increments in *Arctica islandica* L. from the North Sea by means of oxygen and carbon isotopes", *Netherlands Journal of Sea Research*, vol. 33, no. 1, pp. 91-101.

Witbaard, R., Duineveld, G. C. A., & De Wilde, P. A. W. J. 1999, "Geographical differences in growth rates of *Arctica islandica* (Mollusca: Bivalvia) from the North Sea and adjacent waters", *Journal of the Marine Biological Association of the United Kingdom*, vol. 79, no. 5, pp. 907-915.

Witbaard, R. & Bergman, M. J. N. 2003, "The distribution and population structure of the bivalve *Arctica islandica* L. in the North Sea: what possible factors are involved?", *Journal of Sea Research*, vol. 50, no. 1, pp. 11-25.

Yates, M. G., Garbutt, R. A., Barratt, D. R., Turk, T., Brown, N. J., Rispin, W. E. R., McGrorty, S., Durell, S. I., Goss-Custard, J., & Murray, E. 2001, Littoral sediments of the Wash and North Norfolk Coast SAC: the 1998 and 1999 surveys of inter-tidal sediment and invertebrates.

Multiple MCZ Regions

- Abella, E. 1998, "Third European Marine Science and Technology Conference, Lisbon, 23-27 May 1998. Project Synopses Volume I: Marine Systems Maerl grounds: habitats of high biodiversity in European Seas," EUR 18220 EN edn, K. G. Barthel et al., eds., European Commission, pp. 169-178.
- Adey, W. H. & Adey, P. J. 1973, "Studies on the biosystematics and ecology of the epilithic crustose Corallinaceae of the British Isles", *British Phycological Journal*, vol. 8, no. 4, pp. 343-407.
- Airoldi, L. & Beck, M. W. 2007, "Loss, status and trends for coastal marine habitats of Europe", *Oceanography and Marine Biology: an annual review*, vol. 45, pp. 345-405.
- Aprahamian, M. & Walker, A. 2008, "Status of eel fisheries, stocks and their management in England and Wales", *Knowl.Managt.Aquatic Ecosyst.* no. 390-391, p. 07.
- Axelsson, M., Dewey, S., Chaddock, S., & Duke, S. 2006, seastar survey, Ocean Quay Marine, Southampton.
- Bamber, R. N. & Irving, P. W. 1994, "Monitoring the growth of *Sabellaria alveolata* in the field", *Memoires du Museum National d'Histoire Naturelle*, vol. 162, no. 0, pp. 612-613.
- Barnes, R. S. K. 1994, "Macrofaunal Community Structure and Life Histories in Coastal Lagoons," in Elsevier Oceanography Series, 60 edn, B. r. Kjerfve, ed., Elsevier, pp. 311-362.
- Barnes, R. S. K. 1999, "What Determines the Distribution of Coastal Hydrobiid Mudsnaills Within North-Western Europe?", *Marine Ecology*, vol. 20, no. 2, pp. 97-110.
- Beck, M. W., Brumbaugh, R. D., Airoldi, L., Carranza, A., Coen, L. D., Crawford, C., Defeo, O., Edgar, G. J., Hancock, B., Kay, M. C., Lenihan, H. S., Luckenbach, M. W., Toropova, C. L., Zhang, G., & Guo, X. 2011, "Oyster Reefs at Risk and Recommendations for Conservation, Restoration, and Management", *BioScience*, vol. 61, no. 2, pp. 107-116.
- Belpaire, C., Geeraerts, C., Evans, D., Ciccotti, E., & Poole, R. 2011, "The European eel quality database: towards a pan-European monitoring of eel quality", *Environmental Monitoring and Assessment*, vol. 183, no. 1-4, pp. 273-284.
- Birchall, R. A. 2007, "Determining The Extent Of *Sabellaria* Reefs During Seabed Surveys", *Offshore Site Investigation And Geotechnics, Confronting New Challenges and Sharing Knowledge*.
- Birchenough, S. & Bremner, J. 2010, shallow and shelf subtidal habitats and ecology. in mcccip annual report card 2010-11, MCCIP Science Review.
- Bishop, M. J. 1976, "*Hydrobia neglecta* Muus in the British Isles", *Journal of molluscan Studies*, vol. 42, pp. 319-326.

- Bolam, S. G. 2011, "Burial survival of benthic macrofauna following deposition of simulated dredged material", *Environmental Monitoring and Assessment*, vol. 181, no. 1, pp. 13-27.
- Boyd, S. E., Cooper, K. M., Limpenny, D. S., Kilbride, R., Rees, H. L., Dearnaley, M. P., Stevenson, J., Meadows, W. J., & Morris, C. D. 2004, assessment of the re-habilitation of the seabed following marine aggregate dredging, CEFAS, Lowestoft.
- Brossard, J. & Duperret, A. A. 2004, "Coastal chalk cliff erosion: Experimental investigation on the role of marine factors," in *Coastal Chalk Cliff Instability*, R. N. D. Mortimore, ed., pp. 109-120.
- Brown, C. J., Hewer, A. J., Meadows, W. J., Limpenny, D. S., Cooper, K. M., Rees, H. L., & Vivian, C. M. G. 2001, mapping of gravel biotopes and an examination of the factors controlling the distribution, type and diversity of their biological communities, Cefas, Lowestoft.
- Buck, A. L. 1997, An inventory of UK estuaries. Eastern England, 5 edn, Joint Nature Conservation Committee, Peterborough.
- Cabioch, L. 1991, "Space and time series data analysis in coastal benthic ecology. An analytical exercise organised within the framework of the COST 647 project on coastal benthic ecology Evolution a long terme d'ecosystemes benthiques cotiers des sediments subtidaux, en relation avec leur variabilite spatiale physique et biologique. Resultats obtenus par le programme COST 647," B. F. Keegan, ed., Commission of the European Communities, pp. 237-264.
- Collie, J. S., Hall, S. J., Kaiser, M. J., & Poiner, I. R. 2000, "A quantitative analysis of fishing impacts on shelf-sea benthos", *Journal of Animal Ecology*, vol. 69, no. 5, pp. 785-798.
- Constantini, M. L. & Rossi, L. 1995, "Role of fungal patchiness on vegetal detritus in the trophic interactions between two brackish detritivores, *Idotea baltica* and *Gammarus insensibilis*", *Hydrobiologia*, vol. 316, no. 2, pp. 117-126.
- Covey, R. 1991, benthic marine ecosystems in great britain: a review of current knowledge. eastern England and eastern channel (mncr coastal sectors 6 and 7), Nature Conservancy Council.
- Covey, R. 1998, "Eastern Channel (Folkestone to Durlston Head) (MNCR Sector 7)," in *Marine Nature Conservation Review. Benthic marine ecosystems of Great Britain and the north-east Atlantic, Coasts and seas of the United Kingdom. MNCR series edn*, K. Hiscock, ed., Joint Nature Conservation Committee, Peterborough, pp. 199-218.
- Covey, R. 1998, "Chapter 6. Eastern England (Bridlington to Folkestone) (MNCR Sector 6)," in *Marine Nature Conservation Review. Benthic marine ecosystems of Great Britain and the north-east Atlantic*, K. Hiscock, ed., Joint Nature Conservation Committee. (Coasts and seas of the United Kingdom. MNCR series.), Peterborough, pp. 179-198.
- Crisp, D. A. 1964, "The effects of the severe winter of 1962-63 on marine life in Britain", *Journal of Animal Ecology*, vol. 33, no. 1, pp. 165-210.

Curtis, J. M. R. 2007, "Validation of a method for estimating realized annual fecundity in a multiple spawner, the long-snouted seahorse (*Hippocampus guttulatus*), using underwater visual census", *Fishery Bulletin*, vol. 105, no. 3, pp. 327-336.

Davies, J. S., Hall-Spencer, J., Howell, K. L., Roberts, J. M., Hughes, D., & Narayanaswamy, B. E. 1900, *The distribution of cold-water corals on UK banks and seamounts* National Oceanography Center, Southampton, UK.

Diesing, M., Vanstaen, K., & Coggan, R. 1900, *Bedrock Geology Governs Benthic Habitat in the Central English Channel* Association of American Geographers, 1710 16th St, NW Washington, DC 20009 USA.

Driver, P. A. 1975, *The crawfish, *Palinurus elephas* Fabr.: its biology and fishery in British waters*, Council for National Academic Awards.

Edwards, E. 1997, NOAA Technical Report NMFS, vol. 0, no. 129, pp. 85-99.

Edwards, J. R. 1987, "The UK heritage coasts: An assessment of the ecological impacts of tourism", *Annals of Tourism Research*, vol. 14, no. 1, pp. 71-87.

Ellis, J. R., Dulvy, N. K., Jennings, S., Parker-Humphreys, M., & Rogers, S. I. 2005, "Assessing the status of demersal elasmobranchs in UK waters: a review", *Journal of the Marine Biological Association of the United Kingdom*, vol. 85, no. 05, pp. 1025-1047.

Ellis, J. R., McCully, S. R., & Brown, M. J. 2012, "An overview of the biology and status of undulate ray *Raja undulata* in the north-east Atlantic Ocean", *Journal of Fish Biology*.

Fowler, S. L. & Tittley, I. 1993, *the marine nature conservation importance of british coastal chalk cliff habitats*, *English Nature*, 32.

Frederick, S. R. & Maurice, Y. A. 1973, "Taxonomic Index," in *Advances in Marine Biology*, Volume 11 edn, Academic Press, pp. 277-287.

Fretter, V. & Graham, A. 1978, "Prosobranch Mollusks Of Britain And Denmark .3. Neritacea, Viviparacea, Valvatacea, Terrestrial And Freshwater Littorinacea And Rissoacea", *Journal of molluscan Studies* pp. 101-150.

Gates, A. R. 2006, *The physiological ecology of the specialist lagoon amphipod, *Gammarus insensibilis**, University of Southampton.

George, J. D. & Fincham, A. A. "Macro-invertebrate communities of chalk shores in Southeastern England", in *Topics in Marine Biology: Proceedings of the 22nd European Marine Biology Symposium*, Scientia Marina (Barcelona), 53(2-3), Barcelona, Spain, pp. 373-385.

Godwin, H. 1943, "Coastal peat beds of the British Isles and North Sea", *Journal of Ecology*, vol. 31, pp. 199-247.

- Hall-Spencer, J. M. & Stehfest, K. 2009, background document for lophelia pertusa reefs.
- Hendrick, V. J., Foster-Smith, R. L., Davies, A. J., & Newell, R. C. 2011, biogenic reefs and the marine aggregate industry, Cefas.
- Hetherington, S., Quayle, V., & Sullivan, K. 2007, programme 15: eastern channel cod, Cefas, Lowestoft.
- Hinojosa, G., Maggs, C. A., & Johnson, M. 2005, "Epifloral And Epifaunal Assemblages Of Natural And Artificially Stabilized Maerl Beds", *Phycologia*, vol. 44, no. 4, p. 45.
- Hiscock, K., Tyler-Walters, H., & Jones, H. 2002, "High level environmental screening study for offshore wind farm developments - marine habitats and species project", Marine Biological Association to the Department of Trade and Industry New & Renewable Energy Programme.
- Hiscock, K. & Oakley, J. 2005, English channel towed sledge seabed images. phase 2: analysis of selected tow images. report to the joint nature conservation committee from the marine biological association. plymouth, Marine Biological Association and JNCC, Plymouth, 90.
- Hiscock, K., Sewell, J., & Oakley, J. 2005, seagrass beds. marine health check 2005. a report to gauge the health of the UK's sea-life, WWF-UK, Godalming.
- Hoeksema, D. F. & Segers, W. A. 1993, "On the systematics and distribution of the marine gastropod *Caecum armoricum* de Folin, 1869 (Prosobranchia, Caecidae)", *Gloria Maris*, vol. 31, pp. 79-88.
- Hoenselaar, H. J. & Hoenselaar, J. 1990, "On The Identification Of Protoconchs Of Some European Caecidae Gastropoda Prosobranchia", *Basteria*, vol. 54, no. 4-6, pp. 167-169.
- Holme, N. A. & Wilson, J. B. 1985, "Faunas associated with longitudinal furrows and sand ribbons in a tide-swept area in the English Channel", *Journal of the Marine Biological Association of the United Kingdom*, vol. 65, no. 4, pp. 1051-1072.
- Hughes, R. G. & Paramor, O. A. L. 2004, "On the loss of saltmarshes in south-east England and methods for their restoration", *Journal of Applied Ecology*, vol. 41, no. 3, pp. 440-448.
- Jennings, S. & Kaiser, M. J. 1998, "The Effects of Fishing on Marine Ecosystems," in *Advances in Marine Biology*, Volume 34 edn, J. H. S. Blaxter, A. J. Southward, & P. A. Tyler, eds., Academic Press, pp. 201-352.
- Künitzer, A., Basford, D., Craeymeersch, J. A., Dewarumez, J. M., D'Arjes, J., Duineveld, G. C. A., Eleftheriou, A., Heip, C., Herman, P., Kingston, P., Niermann, U., Rachor, E., Rumohr, H., & de Wilde, P. A. J. 1992, "The benthic infauna of the North Sea: species distribution and assemblages", *ICES Journal of Marine Science: Journal du Conseil*, vol. 49, no. 2, pp. 127-143.

Kennedy, C. R. & Guégan, J. F. 1996, "The number of niches in intestinal helminth communities of *Anguilla anguilla*: are there enough spaces for parasites?", *Parasitology*, vol. 113, no. 03, pp. 293-302.

Kenny, A. J., Rees, H. L., & Lees, R. G. 1991, "An inter-regional comparison of gravel assemblages off the English east and south coasts: Preliminary results", ICES, Copenhagen(Denmark).1991.

Kenyon, N. H. & Stride, A. H. 1970, "The Tide-Swept Continental Shelf Sediments Between The Shetland Isles And France", *Sedimentology*, vol. 14, no. 3-4, pp. 159-173.

Kenyon, N. H. 1970, "Sand ribbons of European tidal seas", *Marine Geology*, vol. 9, no. 1, pp. 25-39.

Kenyon, N. H., Belderson, R. H., Stride, A. H., & Johnson, M. A. 2009, "Offshore Tidal Sand-Banks as Indicators of Net Sand Transport and as Potential Deposits," in *Holocene Marine Sedimentation in the North Sea Basin*, Blackwell Publishing Ltd., pp. 257-268.

Killeen, I. J. & Light, J. M. 2000, "Sabellaria, a polychaete host for the gastropods *Noemiamea dolioliformis* and *Graphis albida*", *Journal of the Marine Biological Association of the UK*, vol. 80, no. 03, pp. 571-573.

Killeen, I. J. & Light, J. M. 2004, "The status, distribution and ecology of *Paludinella littorina* (delle Chiaje, 1828) (Gastropoda: Assimineidae) in the British Isles", *Newsletter.Porcupine Marine Natural History Society*, vol. 16, p. 36.

Langston, W. J., Chesman, B. S., Burt, G. R., Pope, N. D., & McEvoy, J. 2002, "Metallothionein in liver of eels *Anguilla anguilla* from the Thames Estuary: an indicator of environmental quality?", *Marine Environmental Research*, vol. 53, no. 3, pp. 263-293.

Martin, C. S., Vaz, S., Ellis, J. R., Coppin, F., Le Roy, D., & Carpentier, A. 2010, "Spatio-temporal patterns in demersal elasmobranchs from trawl surveys in the eastern English Channel (1988-2008)", *Marine Ecology Progress Series*, vol. 417, pp. 211-228.

McBreen, F., Wilson, J., Mackie, A., & Nic Aonghusa, C. o. 2008, "Seabed mapping in the southern Irish Sea: predicting benthic biological communities based on sediment characteristics", *Hydrobiologia*, vol. 606, no. 1, pp. 93-103.

Mieszkowska, N., Kendall, M., Hawkins, S., Leaper, R., Williamson, P., Hardman-Mountford, N., & Southward, A. 2006, "Changes in the range of some common rocky shore species In Britain - A Response To Climate Change?", *Marine Biodiversity* pp. 241-251.

Milner, N. 2001, "At the Cutting Edge: Using Thin Sectioning to Determine Season of Death of the European Oyster, *Ostrea edulis*", *Journal of Archaeological Science*, vol. 28, no. 8, pp. 861-873.

Morgan, T. S., Rogers, A. D., & Iyengar, A. 2000, "Novel microsatellite markers for the European oyster *Ostrea edulis*", *Molecular Ecology*, vol. 9, no. 4, pp. 495-497.

Oakley, J. & Hiscock, K. 2005, English channel towed sledge seabed images. phase 1: scoping study and example analysis. report to the joint nature conservation committee from the marine biological association, The Marine Biological Association of the UK.

Palero, F., Abell³, P., Macpherson, E., Matthee, C. A., & Pascual, M. A. 2010, "Genetic diversity levels in fishery-exploited spiny lobsters of the genus *Palinurus* (Decapoda: Achelata)", *Journal of Crustacean Biology*, vol. 30, no. 4, pp. 658-663.

Paul, A. R. 1958, "Eastbourne Marine Shore Fauna: Being notes on the distribution of species occurring in the neighbourhood of the Beachy Head and Holywell areas of Eastbourne", *Proceedings of the Zoological Society of London*, vol. 131, no. 4, pp. 527-557.

Paul, M., Lefebvre, A., Manca, E., & Amos, C. L. 2011, "An acoustic method for the remote measurement of seagrass metrics", *Estuarine, Coastal and Shelf Science*.

Pearce, B., Tappin, D. R., Dove, D., & Pinnion, J. 2012, "42 - Benthos Supported by the Tunnel-Valleys of the Southern North Sea," in *Seafloor Geomorphology as Benthic Habitat*, Elsevier, London AB - *Sabellaria Alveolata* Reefs, pp. 597-612.

Pearson, C. 2003, The genetic population structure of the lagoon specialists *Nematostella vectensis*, *Cerastoderma glaucum* and *Gammarus insensibilis* from populations along the southern and eastern coasts of the United Kingdom, University of Southampton.

Pearson, C. V. M., Rogers, A. D., & Shearer, M. 2002, "The genetic structure of the rare lagoonal sea anemone, *Nematostella vectensis* Stephenson (Cnidaria: Anthozoa) in the United Kingdom based on RAPD analysis", *Molecular Ecology*, vol. 11, no. 11, pp. 2285-2293.

Posford, D. E. 1999, overview of coastal sand dunes, saltmarsh and vegetated shingle by natural area 317.

Potts, G. W. & Swaby, S. E. 1992, The current status of the giant goby, *Gobius cobitis pallas*, in the British Isles, Unpublished contract report no. 99 F2A 059 edn, Joint Nature Conservation Committee, Peterborough AB - *G. cobitis*.

Price, J. H., Tittley, I., & Honey, S. I. 1977, "The benthic marine flora of Lincolnshire and Cambridgeshire: a preliminary survey. Part 2", *Naturalist*, vol. 102, pp. 91-104.

Price, J. H., Tittley, I., & Richardson, W. D. 1979, "The distribution of *Padina pavonica* (L.) Lamour. (Phaeophyta: Dictyotales) on British and adjacent European shores", *Bulletin of the British Museum (Natural History), Botany*, vol. 7, pp. 1-67.

Price, J. H. 1984, "The distribution of benthic marine algae a bibliography for the British Isles Supplement 4", *British Phycological Journal*, vol. 19, no. 4, pp. 385-404.

Quero, J. C. & Gueguen, J. 1980, "The Rays of the Celtic Sea and the Bristol Channel. Abundance and Distribution", *SCI.PECHE*. no. 318, pp. 1-22.

- Ranwell, D. S., Wyer, D. W., Boorman, L. A., Pizzey, J. M., & Waters, R. J. 1974, "Zostera transplants in Norfolk and Suffolk, Great Britain", *Aquaculture*, vol. 4, no. 0, pp. 185-198.
- Rees, E. I. S., Bergmann, M., Galanidi, M., Hinz, H., Shucksmith, R., & Kaiser, M. J. 2005, "An enriched Chaetopterus tube mat biotope in the eastern English Channel", *Journal of the Marine Biological Association of the UK*, vol. 85, no. 2, pp. 323-326.
- Reiss, H., Degraer, S., Duineveld, G. C. A., KrÄncke, I., Aldridge, J., Craeymeersch, J. A., Eggleton, J. D., Hillewaert, H., Lavaleye, M. S. S., Moll, A., Pohlmann, T., Rachor, E., Robertson, M., Vanden Berghe, E., van Hoey, G., & Rees, H. L. 2010, "Spatial patterns of infauna, epifauna, and demersal fish communities in the North Sea", *ICES Journal of Marine Science: Journal du Conseil*, vol. 67, no. 2, pp. 278-293.
- Rogers, S. I., Millner, R. S., & Mead, T. A. 1998, the distribution and abundance of young fish on the east and south coast of England (1981 to 1997), CEFAS, Lowestoft, 108.
- Rogers, S. I., Maxwell, D., Rijnsdorp, A. D., Damm, U., & Vanhee, W. 1999, "Fishing effects in northeast Atlantic shelf seas: patterns in fishing effort, diversity and community structure. IV. Can comparisons of species diversity be used to assess human impacts on demersal fish faunas?", *Fisheries Research*, vol. 40, no. 2, pp. 135-152.
- Rogers, S. I. & Ellis, J. R. 2000, "Changes in the demersal fish assemblages of British coastal waters during the 20th century", *ICES Journal of Marine Science: Journal du Conseil*, vol. 57, no. 4, pp. 866-881.
- Rowland, S. J., Donkin, P., Smith, E. L., Widdows, J., & Lowe, D. M. 2002, characterisation of bioaccumulated and unidentified agents causing reduced scope for growth in the blue mussel (*mytilus edulis*, University of Plymouth, Plymouth (UK).
- Rumohr, H. & Krost, P. 1991, "Experimental evidence of damage to benthos by bottom trawling with special reference to *Arctica islandica*", *Meeresforschung*, vol. 33, no. 4, pp. 340-345.
- Seaward, D. R. 1989, "Caecum-Armoricum (Prosobranchia, Rissoacea) New To The British Marine Fauna", *Journal of Conchology*, vol. 33, p. 268.
- Stride, A. H., Wilson, J. B., & Curry, D. 1999, "Accumulation of late Pleistocene and Holocene biogenic sands and gravels on the continental shelf between northern Scotland and western France", *Marine Geology*, vol. 159, no. 1, pp. 273-285.
- Swaby, S. E. & Potts, G. W. 1995, The current status and distribution of the giant goby (*Gobius cobitis*) in the British Isles *English Nature*, Peterborough AB - *G. cobitis*.
- Tesch, F. W. 1980, "Occurrence of eel, *Anguilla anguilla*, larvae west of the European continental shelf, 1971-1977", *Environmental Biology of Fishes*, vol. 5, pp. 185-190.

Thomson, A. G., Fuller, R. M., Yates, M. G., Brown, S. L., Cox, R., & Wadsworth, R. A. 2003, "The use of airborne remote sensing for extensive mapping of intertidal sediments and saltmarshes in eastern England", *International Journal of Remote Sensing*, vol. 24, no. 13, pp. 2717-2737.

Tittley, I. 1982, "The effects of man-made constructions on the marine flora of the southern North Sea", The Open University.

Tittley, I., Spurrier, C. J. H., Chimonides, P. J., George, J. D., Moore, J. A., Evans, N. J., & Muir, A. I. 1998, "Survey of chalk cave, cliff, intertidal and subtidal reef biotopes in the Thanet coast cask", *English Nature Research Reports*, vol. 325, 100p..

Tittley, I., Gilliland, P., & Pound, D. 1998, "Changes in the marine flora of the North Sea. Marine flora of the Thanet coast marine SAC: A conservation and management perspective," G. W. Scott & I. Tittley, eds., *Centre for Environmental Research into Coastal Issues (CERCI)*, Scarborough, pp. 65-75.

Van Hoey, G., Guilini, K., Rabaut, M., Vincx, M., & Degraer, S. 2008, "Ecological implications of the presence of the tube-building polychaete *Lanice conchilega* on soft-bottom benthic ecosystems", *Marine Biology*, vol. 154, no. 6, pp. 1009-1019.

Warwick, R. M. & Somerfield, P. J. 2010, "The structure and functioning of the benthic macrofauna of the Bristol Channel and Severn Estuary, with predicted effects of a tidal barrage", *Marine Pollution Bulletin*, vol. 61, no. 1, pp. 92-99.

Wharfe, J. R., Wilson, S. R., & Dines, R. A. 1984, "Observations On The Fish Populations Of An East Coast Estuary", *Marine Pollution Bulletin*, vol. 15, no. 4, pp. 133-136.

Williams, R. A. 1975, "A redescription of the brackish-water sea anemone *Nematostella vectensis* Stephenson, with an appraisal of congeneric species", *Journal of Natural History*, vol. 9, no. 1, pp. 51-64.

Witbaard, R. & Klein, R. 1994, "Long-term trends on the effects of the southern North Sea beamtrawl fishery on the bivalve mollusc *Arctica islandica* L. (Mollusca, bivalvia)", *ICES Journal of Marine Science*, vol. 51, no. 1, pp. 99-105.

Wolters, M., Bakker, J. P., Bertness, M. D., Jefferies, R. L., & Moller, I. 2005, "Saltmarsh erosion and restoration in south-east England: squeezing the evidence requires realignment", *Journal of Applied Ecology*, vol. 42, no. 5, pp. 844-851.