

References

- Abeledo, L.G., Calderini, D.F., Slafer, G.A. (2008). Nitrogen economy in old and modern malting barleys. *Field Crops Research* 106, 171-178.
- Adamsen, F.J., Pinter Jr., P.J., Barnes, E.M., LaMorte, R.L., Wall, G.W., Leavitt, S.W., Kimball, B.A. (1999). Measuring wheat senescence with a digital camera. *Crop Science* 39, 719-724.
- Anon (2000). Fertiliser Recommendations for Agricultural and Horticultural crops. (Seventh Edition). *MAFF (now DEFRA) Reference Book 209*. London: HMSO.
- Anon (2006). De Klein, C.A.M., Novoa, R.S.A., Ogle, S.M., Smith, K.A., Rochette, P., Wirth, T.C., McConkey, B.G., Mosier, A., Rypdal, K. Chapter 11: N₂O Emissions from managed soils, and CO₂ emissions from lime and urea application. In: *2006 IPCC Guidelines for National Greenhouse Gas Inventories; Volume 4: Agriculture, Forestry and Other Land Use*: International Panel on Climate Change.
- Anon (2007). Project NF0614 'Environmental Assessment Tool for Biomaterials' Funded by Defra under Programme CTD0402 'Supply Chain Assessment and Development for Industrial Materials from Crops'. ([www.nnfcc.co.uk<Toolkits<Assessment tools](http://www.nnfcc.co.uk/Toolkits/Assessment%20tools)).
- Anon (2010). Fertiliser Recommendations for Agricultural and Horticultural Crops (RB209, 8th edition). *Defra, London (expected to become available from The Stationery Office, Norwich in 2010; near-final drafts were used here)*.
- Arcadia Biosciences (2007). <http://www.arcadiabio.com/assets/media/71010-ACPFGCSIRO.pdf>
- Austin, R.B., Bingham, J., Blackwell, R.D., Evans, L.T., Ford, M.A., Morgan, C.L., Taylor, M. (1980). Genetic improvements in winter wheat yields since 1900 and associated physiological changes. *Journal of Agricultural Science* 94, 675-689.
- Babar, M.A., Reynolds, M.P., van Ginkel, M., Klatt, A.R. Raun, W.R., Stone, M.L. (2006). Spectral reflectance indices as a potential indirect selection criteria for wheat yield under irrigation. *Crop Science* 46, 578–588.
- Behrens, T., Kraft, M., Wiesler, F. (2004). Influence of measuring angle, nitrogen fertilization, and variety on spectral reflectance of winter oilseed rape canopies. *Journal of Plant Nutrition and Soil Science* 167, 99-105.
- Berry, P., Foulkes, Spink, J., Teakle, G., White, P. (2008). Breeding for improved nitrogen use efficiency in oilseed rape. In: Resource capture by crops: integrated approaches. AAB 10-12 September 2008, Sutton Bonington, University of Nottingham, UK.
- Berry, P.M., Kindred, D.R., Paveley, N.D. (2008). Quantifying the effects of fungicides and disease resistance on greenhouse gas emissions associated with wheat production. *Plant Pathology* 57, 1000–1008.
- Berry, P.M., Spink, J., Foulkes, J., White, P.J. (2010). The physiological basis of genotypic differences in nitrogen use efficiency in oilseed rape (*Brassica napus* L.). *In press*
- Berry, P.M. and Spink, J.H. (2009). 'Canopy Management' and late nitrogen applications to improve yield of oilseed rape. *Home-Grown Cereals Authority Project Report No 447*. HGCA, London, 212 pp.

- Brentrup, F. and Paliere, C. (2008). GHG emissions and energy efficiency in european nitrogen fertiliser production and use. In: Proceedings of The International Fertiliser Society Conference, 11th December 2008. Cambridge, UK.
- Brown, L., Syed, B., Jarvis, S.C., Sneath, R.W., Phillips, V.R., Goulding, K.W.T., Li, C. (2002). Development and application of a mechanistic model to estimate emission of nitrous oxide from UK agriculture. *Atmospheric Environment* 36, 917-928.
- Burton, D.L., Zebarth, B.J., Gillarn, K.M., and MacLeod, J.A. (2008). Effect of split application of fertilizer nitrogen on N₂O emissions from potatoes. *Canadian Journal of Soil Science* 88, 229-239.
- Clarke, S., Kindred, D., Weightman, R. M., Sylvester-Bradley, R. (2008). Growing Wheat for Alcohol and Bioethanol Production in the North East: *An ADAS report commissioned by One North East and North East Processing Industries Cluster*.
- Clayton H., McTaggart I.P., Parker J., Swan L. and Smith K.A. (1997). Nitrous oxide emissions from fertilised grassland: A 2-year study of the effects of N fertiliser form and environmental conditions. *Biology and Fertility of Soils* 25, 252-260.
- Colenne, C., Meynard, J.M., Roche, R., Reau, R. (2002). Effects of nitrogen deficiencies on autumnal growth of oilseed rape. *European Journal of Agronomy* 17, 11-28.
- Chantigny, M.H., Prevost, D., Angers, D.A., Simard, R.R., and Chalifour, F.P. (1998). Nitrous oxide production in soils cropped to corn with varying N fertilization. *Canadian Journal of Soil Science* 78, 589-596.
- Dampney, P.M.R., Salmon, S.E., Greenwell, P., Pritchard, P.E. (1995). Management of breadmaking wheat: Effects of extra nitrogen on yield, grain and flour quality. *Home-Grown Cereals Authority Project Report No 109. HGCA, London*.
- Dampney, P.M.R., Dyer, C.J., Goodlass, G. and Chambers, B. (2006) WP1a – Crop responses to different N fertiliser materials. *Component report for Defra Project NT2605 (CSA 6579) 110pp. Defra, London*.
- Dampney, P.M.R., Edwards, A., Dyer, C.J. (2007). Managing nitrogen applications to new Group 1 and 2 wheat varieties. *Home-Grown Cereals Authority Project Report No 400. HGCA, London*.
- Davidson E A (1991) Fluxes of nitrous oxide and nitric oxide from terrestrial ecosystems. *In* Microbial production and consumption of greenhouse gases: methane, nitrogen oxides and halomethanes, Eds J E Rogers and W B Whitman. pp 219-235. American Society of Microbiology, Washington D.C.
- Defra (2005). A study of the scope for the application of crop genomics and breeding to increase nitrogen economy within cereal and rapeseed based food chains. *Project Report AR0714. Defra, London*.
- Defra (2007) The British survey of fertiliser practice: Fertiliser use on farm crops for crop year 2006. <https://statistics.defra.gov.uk/esg/bsfp.htm>. Accessed 02/03/2010.
- Defra (2008) The British survey of fertiliser practice: Fertiliser use on farm crops for crop year 2007. <https://statistics.defra.gov.uk/esg/bsfp.htm>. Accessed 02/03/2010.
- Defra (2009). The British survey of fertiliser practice: Fertiliser use on farm crops for crop year 2008. <https://statistics.defra.gov.uk/esg/bsfp.htm>. Accessed 02/03/2010.

- Defra (2010). United Kingdom cereal yields 1885 onwards.
https://statistics.defra.gov.uk/esg/datasets/cyield_c.xls. Accessed 15/03/2010.
- del Grosso, S.J., Ojima, D.S., Parton, W.J., and Mosier, A.R. (2002). Simulated effects of tillage and timing of N fertilizer application on net greenhouse gas fluxes and N losses for agricultural soils in the Midwestern USA. In "*Non-CO₂ Greenhouse Gases: Scientific Understanding, Control Options and Policy Aspects*" (J. VanHam, A.P.M. Baede, R. Guicherit and J.G.F. WilliamsJacobse, eds.), pp. 23-28. Millpress Science Publishers, Rotterdam, NL.
- del Prado, A., Brown, L., and Scholefield, D. (2004). NGAUGE DSS as a tool to assist UK dairy farmers to comply with EU nitrate legislation. In "*Controlling Nitrogen Flows and Losses*" (D. J. Hatch, D. R. Chadwick, S. C. Jarvis and J. A. Roker, eds.), pp. 423-424. Wageningen Academic Publishers.
- Dobbie K.E., McTaggart I.P. and Smith K.A. (1999). Nitrous oxide emissions from intensive agricultural systems: Variations between crops and seasons, key driving variables, and mean emissions factors. *Journal of Geophysical Research* 104, 26,891-26,899.
- Dobbie K.E. and Smith K.A. (2001). The effects of temperature, water-filled pore space and land use on N₂O emissions from an imperfectly drained gleysol. *European Journal of Soil Science* 52, 667-673.
- Dobbie K.E. and Smith K.A. (2003). Nitrous oxide emission factors for agricultural soils in Great Britain: the impact of soil water-filled pore space and other controlling variables. *Global Change Biology* 9, 204-218.
- Dhugga, K. S., and Waines, J. G. (1989). Analysis of nitrogen accumulation and use in bread and durum wheat. *Crop Sci.* 29, 1232–1239.
- Drew, M.C., Saker, L.R., Ashley, T.W. (1973). Nutrient supply and the growth of the seminal root system in barley: I. The effect of nitrate concentration on the growth of axes and laterals. *Journal of Experimental Botany* 24, 1189-1202.
- Etter, L. (2007). Power plant in china; a plan to turn rice into carbon credits. *Wall Street Journal*, 9 October 2007.
<http://online.wsj.com/public/article/SB119187524509952568.html>
- Farrer, D.C., Weisz, R., Heiniger, R., Murphy, J.P., White, J.G. (1998). Minimizing Protein Variability in Soft Red Winter Wheat: Impact of Nitrogen Application Timing and Rate. *Agronomy Journal* 98, 1137-1145.
- Firestone M K and Davidson E A (1989) Microbiological basis of NO and N₂O production and consumption in soil. In *Exchange of Trace Gases between Terrestrial Ecosystems and the Atmosphere*, Eds M O Andreae and D S Schimel. pp 7-21. John Wiley & Sons, Inc., New York.
- Foulkes, M.J., Sylvester-Bradley, R., Weightman, R., Snape, J.W. (2007). Identifying physiological traits associated with improved drought resistance in winter wheat. *Field Crops Research* 103, 11-24.
- Foulkes, M.J., Hawkesford, M.J., Barraclough, P.B., Holdsworth, M.J., Kerr, S., Kightley, S., Shewry, P.R. (2009). Identifying traits to improve the nitrogen economy of wheat: Recent advances and future prospects. *Field Crops Research* 114, 329-342.
- George, B.J. (1984). Design and interpretation of nitrogen response experiments. In 'The nitrogen requirement of cereals', *Reference Book 385*, 133-149, *MAFF*.

- Good A.G., Shrawat A.K., Muench D.G. (2004) Can less yield more? Is reducing nutrient input into the environment compatible with maintaining crop production? *Trends in Plant Science* 9, 597–605.
- Good AG, Johnson SJ, De Pauw M, Carroll RT, Savidov N, Vidmar J, Lu Z, Taylor G & Stroehrer V. (2007). Engineering nitrogen use efficiency with alanine aminotransferase. *Canadian Journal of Botany* 85, 252-262.
- Górny, A.G., Ratajczak, D. (2008). Efficiency of nitrogen and phosphorus utilization in progenies of factorial crosses between European and exotic cultivars of spring barley (*Hordeum vulgare* L.). *Journal of Applied Genetics* 49, 349-355.
- Grant, C.A., Brown, K.R., Racz, G.J., Bailey, L.D. (2002). Influence of source, timing and placement of nitrogen fertilization on seed yield and nitrogen accumulation in the seed of canola under reduced-and conventional-tillage management. *Canadian Journal of Plant Science* 82, 629-638.
- Guarda, G., Padovan, S., Delogu, G. (2004). Grain yield, nitrogen-use efficiency and baking quality of old and modern Italian bread-wheat cultivars grown at different nitrogen levels. *European Journal of Agronomy* 21, 181-192.
- Hodge, A. (2004). The plastic plant: Root responses to heterogeneous supplies of nutrients. *New Phytologist* 162, 9-24.
- Howard, D.D., Newman, M.A., Essington, M.E., Percell, W.M. (2002). Nitrogen fertilization of conservation-tilled wheat. II. Timing of application of two nitrogen sources. *Journal of Plant Nutrition* 25, 1329-1339.
- IPCC (1997) Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. Houghton, J.T., Meira Filho, L.G., Lim, B., Tréanton, K., Mamaty, I., Bonduki, Y., Griggs, D.J. and Callander, B.A. (Eds). Intergovernmental Panel on Climate Change (IPCC), IPCC/OECD/IEA, Paris, France.
- Joshi, A.K., Kumari, M., Singh, V.P., Reddy, C.M., Kumar, S., Rane, J., Chand, R. (2007). Stay green trait: Variation, inheritance and its association with spot blotch resistance in spring wheat (*Triticum aestivum* L.). *Euphytica* 153, 59-71.
- Kamh, M., Wiesler, F., Ulas, A., Horst, W.J. (2005). Root growth and N-uptake activity of oilseed rape (*Brassica napus* L.) cultivars differing in nitrogen efficiency. *Journal of Plant Nutrition and Soil Science* 168, 130-137.
- Kindred D, Berry P, Burch O, Sylvester-Bradley R. 2008. Effect of nitrogen fertiliser use on greenhouse gas emissions and land use change. *Aspects of Applied Biology* 88, *Effects of climate change on plants: implications for agriculture*, pp. 53–56. Wellesbourne, UK: Association of Applied Biologists.
- Kindred, D.R., Hatley, D., Green, D., Dyer, C., Sylvester-Bradley, R. (2009) Nitrogen timings to wheat crops to optimise Bioethanol Production in the North East. www.adas.co.uk/LinkClick.aspx?fileticket=d_m58J-5eHw%3d&tabid=267
- King, J., Gay, A., Sylvester-Bradley, R., Bingham, I., Foulkes, J., Gregory, P., Robinson, D. (2003). Modelling cereal root systems for water and nitrogen capture: Towards an economic optimum. *Annals of Botany* 91, 383-390.
- Lázzari, M.A., Landriscini, M.R., Echaguè, M.E. (2005). Soil and fertilizer nitrogen uptake patterns in malting barley with near seeding fertilization. *Ciencia del Suelo* 23, 69-77.

- Le Gouis, J., Beghin, D., Heumez, E., Pluchard, P. (2000). Genetic differences for nitrogen uptake and nitrogen utilisation efficiencies in winter wheat. *European Journal of Agronomy* 12, 163-173.
- Lea, P.J., Azevedo, R.A. (2006). Nitrogen use efficiency. 1. The uptake of nitrogen from the soil. *Annals of Applied Biology* 149, 243-247.
- Limaux, F., Recous, S., Meynard, J.-M., Guckert, A. (1999). Relationship between rate of crop growth at date of fertiliser N application and fate of fertiliser N applied to winter wheat. *Plant and Soil* 214, 49-59.
- Lord, E.I., Vaughan, J. (1987). Optimising nitrogen applications for the production of malting barley. *Aspects of Applied Biology* 15, 319-335.
- Lunn, G.D., Spink, J., Stokes, D.T., Wade, A., Clare, R.W., Scott, R.K. (2001). Canopy Management in winter oilseed rape. *Home-Grown Cereals Authority Project Report No. OS49, pp. 86. HGCA, London.*
- Lunn, G.D., Spink, J., Wade, A., Clare, R.W. (2003). Spring remedial treatments to improve canopy structure and yield in oilseed rape. *Home-Grown Cereals Authority Project Report No. OS64, pp. 97. HGCA, London.*
- MacCarthy, J., Thomas, J., Choudrie, S., Passant, N., Thistlethwaite, G., Murrells, T., Watterson, J., Cardenas, L., & Thomson, A. (2010). UK Greenhouse Gas Inventory, 1990-2008: Annual Report for submission under the Framework Convention on Climate Change. AEA Technology plc, Didcot, UK, April 2010.
- Mahmuti, M., West, J.S., Watts, J., Gladders, P., Fitt, B.D.L. 2009. Controlling crop disease contributes to both food security and climate change mitigation. *International Journal of Agricultural Sustainability* 7, 189-202.
- Maidl, F.X., Panse, A., Dennert, J., Ruser, R., Fischbeck, G. (1996). Effect of varied N rates and N timings on yield, N uptake and fertilizer N use efficiency of a six-row and a two-row winter barley. *European Journal of Agronomy* 5, 247-257.
- McTaggart, I.P., Smith, K.A. (1995). The effect of rate, form and timing of fertilizer N on nitrogen uptake and grain N content in spring malting barley. *Journal of Agricultural Science* 125, 341-353.
- Mistele, B., Schmidhalter, U. (2008). Estimating the nitrogen nutrition index using spectral canopy reflectance measurements. *European Journal of Agronomy* 29, 184-190.
- Muller, K., Böttcher, U., Meyer-Schatz, F., Kage, H. (2008). Analysis of vegetation indices derived from hyperspectral reflection measurements for estimating crop canopy parameters of oilseed rape (*Brassica napus* L.). *Biosystems Engineering* 101, 172-
- Ortiz-Monasterio R., J.I., Sayre, K.D., Rajaram, S., McMahon, M. (1997). Genetic progress in wheat yield and nitrogen use efficiency under four nitrogen rates. *Crop Science* 37, 898-904.
- Overthrow, R. (2005). Nitrogen management in spring malting barley for optimum yield and quality. *Home-Grown Cereals Authority Project Report No 367. HGCA, London.*
- Paponov, I.A., Lebedinskai, S., Koshkin, E.I. (1999). Growth analysis of solution culture-grown winter rye, wheat and triticale at different relative rates of nitrogen supply. *Annals of Botany* 84, 467-473.

- Petersen, C.T., Jensen, C.R., Mogensen, V.O. (2002). Analysis of variation of spectral vegetation index measured in differently fertilized field barley. *Communications in Soil Science and Plant Analysis* 33, 1485-1506.
- Petersen, J. (2004). Crop uptake of ¹⁵N labelled fertilizer in spring wheat affected by application time. *Acta Agriculturae Scandinavica Section B: Soil and Plant Science* 54, 83-90.
- Pillinger, C., Paveley, N., Foulkes, M.J., Spink, J. (2005). Explaining variation in the effects of take-all (*Gaeumannomyces graminis* var. *triticici*) on nitrogen and water uptake by winter wheat. *Plant Pathology* 54, 491-501.
- Post, W.M., Kwon, K.C., 2000. Soil carbon sequestration and land-use change: processes and potential. *Global Change Biology* 6, 317-328.
- Rathke, G.W., Behrens, T., Diepenbrock, W. (2008). Integrated nitrogen management strategies to improve seed yield, oil content and nitrogen efficiency of winter oilseed rape (*Brassica napus* L.): A review. *Agriculture, Ecosystems and Environment* 117, 80-108.
- Roberts, R.K., Walters, J.T., Larson, J.A., English, B.C., Howard, D.D. (2004). Effects of disease, nitrogen source, and risk on optimal nitrogen fertilization timing in winter wheat production. *Agronomy Journal* 96, 792-799.
- Rossato, L., MacDuff, J.H., Laine, P., Le Deunff, E., Ourry, A. (2002). Nitrogen storage and remobilisation in *Brassica napus* L. during the growth cycle: effects of methyl jasmonate on nitrate uptake, senescence, growth and VSP accumulation. *Journal of Experimental Botany* 53, 1131-1141.
- Sardana, V., Zhang, G.-P. (2005). Effect of time of nitrogen application on the growth and yield of two barley (*Hordeum vulgare* L.) Cultivars. *Cereal Research Communications* 33, 785-791.
- Searchinger T, Heimlich R, Houghton RA, Dong F, Elobeid A, Fabiosa J, Tokgoz S, Hayes D, Yu T, (2008). Use of U.S. croplands for biofuels increases greenhouse gases through emissions from land-use change. *Science Express* 319, 1238-1240.
- Semenov, M.A., Jamieson, P.D., Martre, P. (2007). Deconvoluting nitrogen use efficiency in wheat: a simulation study. *European Journal of Agronomy*, 26, 283-294.
- Schils, R.L.M., van Groenigen, J.W., Velthof, G.L. and Kuikman, P.J. (2008). Nitrous oxide emissions from multiple combined applications of fertiliser and cattle slurry to grassland. *Plant and Soil* 310: 89-101.
- Siddique K.H.M., Belford R.K., Tennant D. (1990). Root:shoot ratios of old and modern, tall and semi-dwarf wheats in a mediterranean environment. *Plant and Soil*, 121, 89-98.
- Sieling, K., Schröder, H., Hanus, H. (1998). Mineral and slurry nitrogen effects on yield, N uptake and apparent N-use efficiency of oilseed rape (*Brassica napus*). *Journal of Agricultural Science* 130, 165-172.
- Silva, S.A., De Carvalho, F.I.F., Caetano, V.D.R., De Oliveira, A.C., De Coimbra, J.L.M., De Vasconcellos, N.J.S., Lorencetti, C. (2000). Genetic basis of stay-green trait in bread wheat. *Journal of New Seeds* 2, 55-68.
- Simmonds, N.W. (1995). The relation between yield and protein in cereal grain. *Journal of the Science of Food and Agriculture* 68, 309-315.

- Sitthaphanit, S., Limpinuntana, V., Toomsan, B., Panchaban, S., and Bell, R.W. (2009). Fertiliser strategies for improved nutrient use efficiency on sandy soils in high rainfall regimes. *Nutrient Cycling in Agroecosystems* **85**, 123-139.
- Stokes, D.T., Sylvester-Bradley, R. Clare, R.W., Scott, R.K., Hopkinson, J. Milford, G.F.J. & Salmon, S.E. (1998). An integrated approach to nitrogen nutrition of winter wheat. *Home-Grown Cereals Authority Project Report No. 159, HGCA, London.*
- Subbarao, G.V., Tomohiro, B., Masahiro, K., Osamu, I., Samejima, H., Wang, H.Y., Pearse, S.J., Gopalakrishnan, S., Nakahara, K., Zakir Hossain, A.K.M., Tsujimoto, H. & Berry, W.L. (2007). Can biological nitrification inhibition (BNI) genes from 15 perennial *Leymus racemosus* (Triticeae) combat nitrification in wheat farming? *Plant & Soil* **299**, 55–64.
- Svečnjak, Z., Rengel, Z. (2006). Nitrogen utilization efficiency in canola cultivars at grain harvest. *Plant and Soil* **283** (1-2), pp. 299-307.
- Sylvester-Bradley, R., Foulkes, J., Reynolds, M. (2005) Future wheat yields: evidence, theory and conjecture. In: *Yields of farmer species: Constraints and opportunities in the 21st century*. Eds. Sylvester-Bradley, R., Wiseman, J. Nottingham University Press, Nottingham.
- Sylvester-Bradley, R. (2009) Nitrogen for winter wheat – management guidelines. *HGCA, Stoneleigh, Warwickshire. 20 pp.*
- Sylvester-Bradley, R., Kindred, D.R., Blake, J., Dyer, C.J. and Sinclair, A. (2008). Optimising fertiliser nitrogen for modern wheat and barley crops. *Home-Grown Cereals Authority Project Report No. 438, HGCA, London. 116 pp.*
- Sylvester-Bradley, R. & Clarke, S. (2009). Using grain N% as a signature for good N use. *Home-Grown Cereals Authority Project Report No. 458, HGCA, London. 61 pp.*
- Sylvester-Bradley, R., Kindred, D.R., Weightman, R., Thomas, W., Swanson, S., Thompson, D., Fuerhelm, D., Creasy, O., Argillier, O., Melichar, J., Brosnan, J., Agu, R., Cowe, I., Hemingway, D., Robinson, D., Wilcox, S. (2010). Genetic reduction of energy use and emissions of nitrogen through cereal production: GREEN grain. *Home-Grown Cereals Authority Project 2979, in press, HGCA, London.*
- Tambussi, E.A., Nogués, S., Ferrio, P., Voltas, J., Araus, J.L. (2004). Does higher yield potential improve barley performance in Mediterranean conditions? A case study. *Field Crops Research* **91**, 149–160.
- Teakle, G., Berry, P., Foulkes, Spink, J., White, P. (2008). Breeding for improved nitrogen use efficiency in oilseed rape. In: *Resource capture by crops: integrated approaches*. Crucifer 2008 8-11 September 2008, Lillehammer, Norway.
- Thorman, R.E., Sagoo, E., Williams, J.R., Chambers, B.J., Chadwick, D.R., Laws J.A. and Yamulki S (2007). The effect of slurry application timings on direct and indirect N₂O emissions from free draining grassland soils. Proceedings of *The 15th Nitrogen Workshop: Towards a Better Efficiency in N Use*, 28-30 May 2007, Lleida, Spain, pp 297-299.
- Van Ginkel, M., Ortiz-Monasterio, I., Trethowan, R., Hernandez, E. (2001). Methodology for selecting segregating populations for improved N-use efficiency in bread wheat. *Euphytica* **119**, 223-230.
- Van Groenigen, J.W., Schils, R.L.M., Velthof, G.L., Kuikman, P.J., Oudendag, D.A., and Oenema, O. (2008). Mitigation strategies for greenhouse gas emissions from animal

production systems: synergy between measuring and modelling at different scales. *Australian Journal of Experimental Agriculture* 48, 46-53.

Van Vuuren, M.M., Robinson, D., Griffiths, B.S. (1996). Nutrient inflow and root proliferation during the exploitation of a temporarily and spatially discrete source of nitrogen in soil. *Plant and Soil* 178, 185-192.

Varga, B., Svečnjak, Z. (2006). The effect of late-season urea spraying on grain yield and quality of winter wheat cultivars under low and high basal nitrogen fertilization. *Field Crops Research* 96, 125-132.

Verma, V., Foulkes, M.J., Worland, A.J., Sylvester-Bradley, R., Caligari, P.D.S., Snape, J.W. (2004). Mapping quantitative trait loci for flag leaf senescence as a yield determinant in winter wheat under optimal and drought-stressed environments. *Euphytica* 135, 255-263.

Wamser, A.F., Mundstock, C.M. (2007). Protein content in barley seeds affected by nitrogen application in different growth stages. *Ciencia Rural* 37, 1571-1576.

White, E.M. and Wilson, F.E.A. (2006). Responses of grain yield, biomass and harvest index and their rates of genetic progress to nitrogen availability in ten winter wheat varieties. *Irish Journal of Agricultural and Food Research* 45, 85-101.

Cited Defra projects:

AC0101 – An improved inventory of greenhouse gases from agriculture.

AC0206 – A review of research to identify best practice for reducing greenhouse gases from agriculture and land management.

AC0213 - Potential for nitrification inhibitors and fertiliser nitrogen application timing strategies to reduce direct and indirect nitrous oxide emissions from UK agriculture.

IF0175 - A critical review of recent policy-relevant research on nitrogen cycling.

NT2605 – Component report for Defra project NT2605: WP2 The effect of N fertiliser forms on nitrous oxide emissions

RMP4950 - UK marginal abatement cost curves for the agriculture and land use, land-use change and forestry sectors out to 2022, with qualitative analysis of options to 2050.

RMP/5142 - Analysis of policy instruments for reducing greenhouse gas emissions from agriculture, forestry and land management.