

APPENDIX 1A.6

BIBLIOGRAPHY

Adamson, H.F., Ross, S.Y. & Gardner, S.M. (2001). The response of upland wet heath to ESA stocking rates: a heft scale approach. In Biodiversity, plant structure and vegetation heterogeneity: interactions with the grazing herbivore, Crianlarich. November 2002.

ADAS (1998). Monitoring of heather condition in the North Peak ESA 1993-1997. MAFF Publications, London.

ADAS (2001). Extensification and Sheep Performance. Unpublished report to Defra.

Ali, H.A.M., Mayes, R.W., Lamb, C.S., Hector, B.L., Verma, A.K. & Orskov, E.R. (2004). The potential of long-chain fatty alcohols and long-chain fatty acids as diet composition markers: development of methods for quantitative analysis and faecal recoveries of these compounds in sheep fed mixed diets. *Journal of Agricultural Science* 142, 71-78.

Allen, D.S. (1995). Factors Influencing Habitat Associations of Upland Passerines, Particularly the Whinchat (*Saxicola rubetra*): the Importance of Scale. DPhil thesis, University of York.

Alonso, I., Hartley, S.E. & Thurlow, M. (2001). Heather-grass competition on Scottish moorlands: Interacting effects of nutrient enrichment and grazing regime. *Journal of Vegetation Science* 12, 249-260.

Amer, P.R., Emmans, G.C. & Simm, G. (1997). Economic values for carcass traits in UK commercial beef cattle. *Livestock Production Science* 51: 267-281.

Anderson, P. & Radford, E. (1994). Changes in vegetation following reduction in grazing pressure on the National Trust Kinder Estate, Peak District, Derbyshire, England. *Biological Conservation* 69, 55-63.

Anderson, P. & Yalden, D.W. (1981). Increased sheep numbers and the loss of heather moorland in the Peak District, England. *Biological conservation* 20, 195-213.

Anderson, P., Tallis, J.H. & Yalden, D.W. (1997). Restoring moorland: Peak District Moorland Management Project. Phase III Report. Peak Park Joint Planning Board, Bakewell, UK.

Andreev, A.V. (1988) Ecological energetics of Palearctic Tetraonidae in relation to chemical composition and digestibility of their winter diets. *Canadian Journal of Zoology* 66, 1382–1388.

Andrusenko, N.N. (1980). K pitaniyu nekotorykh maibolee massovykh vidov kulikov na ozere Tengiz. *Novoe v izuchenii biologii i rasprostranении kulikov*. Moscow: Nauka.

Anon. (1983). Osnovnye osobennosti pishchevykh svyazei kulikov v basseine r.Shchuch'ei. Moscow: Nauka.

Anon. (1995). Biodiversity, the UK Action Plan. HMSO, London.

Anon. (1999). The White Moorland Programme. In *The Heather Trust Annual Report 1999* (eds J. Phillips & W. Gruellich), pp. 9-11. The Heather Trust, Kippen Stirlingshire.

Anon. (2000). The White Moorland Programme. In *The Heather Trust Annual Report 2000* (eds J. Phillips & W. Gruellich), pp. 8. The Heather Trust, Kippen, Stirlingshire.

Armstrong, H.M., Gordon, I.J., Grant, S.A., Hutchings, N.J., Milne, J.A. & Sibbald, A.R. (1997). A model of the grazing of hill vegetation by sheep in the UK. I. The prediction of vegetation biomass. *Journal of Applied Ecology* 34, 166-185.

Armstrong, H.M., Gordon, I.J., Hutchings, N.J., Illius, A.W., Milne, J.A. & Sibbald, A.R. (1997). A model of the grazing of hill vegetation by sheep in the UK. II. The prediction of offtake by sheep. *Journal of Applied Ecology* 34, 186-207.

Armstrong, R.H., Common, T.G. & Smith, H.K. (1986). The voluntary intake and in vivo digestibility of herbage harvested from indigenous hill plant communities. *Grass and Forage Science* 41, 53-60.

Arnold, G.W. (1960). The effect of the quantity and quality of pasture available to sheep on their grazing behaviour. *Australian Journal of Agricultural Research* 11, 1026-1033.

Arnold, G.W. (1984). Comparison of the time budgets and circadian patterns of maintenance activities in sheep, cattle and horses grouped together. *Applied Animal Behaviour Science* 13, 19-30.

Arnold, G.W., Ball, J., Mcmanus, W.R. & Bush, I.R. (1966). Studies on the diet of the grazing animal. I Seasonal changes in the diet of sheep grazing on pastures of

different availability and composition. *Australian Journal of Agricultural Research* 17, 543-556.

Artem'yev, Y.T. & Popov, V.A. (1978). *Ptitsy Volzhsko-Kamskogo Kraya*. Moscow: Nauka.

Backshall, J., Manley, J. & Rebane, M.E. (2001). *The Upland Management Handbook*. English Nature, Peterborough, UK.

Baines, D. (1990). The roles of predation, food and agricultural practice in determining the breeding success of lapwing *Vanellus vanellus* on upland grasslands. *Journal of Animal Ecology* 59, 915-929.

Baines, D. (1996). The implications of grazing and predator management on the habitats and breeding success of black grouse *Tetrao tetrix*. *Journal of Applied Ecology* 33, 54-62.

Baker, M.C. (1977). Shorebird food habits in the eastern Canadian Arctic. *Condor* 79, 56-62.

Bardgett, R.D., Marsden, J.H. & Howard, D.C. (1995). The extent and condition of heather on moorland in the uplands of England and Wales. *Biological Conservation* 71, 155-161.

Bardgett, R.D., Marsden, J.H., Howard, D.C. & Hossell, J.E. (1995). The extent and condition of heather in moorland, and the potential impact of climate change. In Thompson, D.B.A., Hester, A.J. & Usher, M.B. (eds) *Heaths and Moorland: Cultural Landscapes*: 43-50. Edinburgh: HMSO.

Bauchop, T. (1981). The anaerobic fungi in fibre digestion. *Agriculture and the Environment* 6, 338 - 348.

Beale, C.M, Monaghan, P. (2004). Human disturbance: people as predation-free predators? *Journal of Applied Ecology* 41, 335-343.

Beintema, A.J., Thissen, J.B., Tensen, D. & Visser, G.H. (1991). Feeding ecology of Charadriiform chicks in agricultural grassland. *Ardea* 79, 31-44.

Bell, J.R., Wheater, C.P. & Cullen, W.R. (2001). The implications of grassland and heathland management for the conservation of spider communities: a review. *Journal of Zoology*, London 255, 377-387.

- Bell, R.H.V. (1970). The use of the herb layer by grazing ungulates in the Serengeti. In: Watson A. (ed). *Animal Populations in Relation to their Food Resources*. pp. 111-123. Oxford: Blackwell Scientific Publications.
- Belovsky, G.E. (1986). Generalist herbivore foraging and its role in competitive interactions. *American Zoologist* 26, 51-69.
- Benton, T.G., Vickery, J.A., Wilson, J.D. (2003). Farmland biodiversity: is habitat heterogeneity the key? *Trends in Ecology and Evolution* 18, 182-188.
- Berry, P.M., Vanhinsberg, D., Viles, H.A., Harrison, P.A., Pearson, R.G., Fuller, R.J., Butt, N. & Miller, F. (2001). Impacts on terrestrial environments. In Harrison, P.A., Berry, P.M. & Dawson, T.P. (eds) *Climate Change and Nature Conservation in Britain and Ireland: Modelling Natural Resource Responses to Climate Change, the MONARCH Project: 43–149*. UKCIP Technical Report. Oxford: UKCIP.
- Binzenhofer, B., Schroeder, B., Strauss, B., Biedermann, R. & Settele, J. (2005). Habitat models and habitat connectivity analysis for butterflies and burnet moths - The example of *Zygaena carniolica* and *Coenonympha arcania* *Biological Conservation* 126, 247-259.
- Birch, C.P.D., Vuichard, N. & Werkman, B.R. (2000). Modelling the effects of patch size on vegetation dynamics: Bracken (*Pteridium aquilinum* (L.) Kuhn) under grazing. *Annals of Botany* 85, 63-76.
- Birch, C.P.D., Werkman, B.R. & Partridge, L.W. (1997). A predictive model of vegetation dynamics under grazing. In: *Proceedings of the 18th International Grassland Conference, Winnipeg, Canada, 1997*. pp. 12:5-12:6.
- Bokdam, J. & Gleichman, J.M. (2000). Effects of grazing by free-ranging cattle on vegetation dynamics in a continental north-west European heathland. *Journal of Applied Ecology* 37, 415-431.
- Bokdam, J. (2001). Effects of browsing and grazing on cyclic succession in nutrient-limited ecosystems. *Journal of Vegetation Science* 12, 875–886.
- Borchtchevski, V.G. (2000). Diet of the Black Grouse *Tetrao tetrix* in Vodlozerski National Park, North-western Russia. *Game and Wildlife Science* 17, 29–42.

- Britshgi, A., Spaar, R. & Arlettaz, R. (2006). Impact of grassland farming intensification on the breeding ecology of an indicator insectivorous passerine, the Whinchat *Saxicola rubetra*: Lessons for overall Alpine meadowland management. *Biological Conservation* 130, 193-205.
- Britton, A. & Marrs, R. (2000). Management techniques to promote *Calluna vulgaris* cover on heathland invaded by grasses. *Aspects of Applied Biology* 58, 173-178.
- Brooke, M. (1981). How an adult wheatear *Oenanthe oenanthe* uses its territory when feeding nestlings. *Journal of Animal Ecology* 50, 683–696.
- Brown D.R. (2000). A review of bio-economic models. Paper prepared for Cornell African Food Security and Natural Resource Management (CAFSNRM) Program, http://aem.cornell.edu/faculty_sites/cbb2/grad/BrownReviewofBioeconomicModels.pdf
- Brown, A.F. & Stillman, R.A. (1993). Bird-habitat associations in the eastern Highlands of Scotland. *Journal of Applied Ecology* 30: 31-42.
- Buchanan, G, Pearce-Higgins, J, Grant, M., Robertson, D. & Waterhouse, A. (2005). Characterization of moorland vegetation and the prediction of bird abundance using remote sensing. *Journal of Biogeography* 32, 697-707.
- Buchanan, G.M., Grant, M.C., Sanderson, R.A. & Pearce-Higgins, J.W. (2006a). The contribution of invertebrate taxa to moorland bird diets and the potential implications of land-use management. *Ibis* 148, 615-628.
- Buchanan, G.M., Pearce-Higgins, J.W. & Grant, M.C. (2006b). Observer variation in estimates of meadow pipit *Anthus pratensis* and skylark *Alauda arvensis* abundance on moorland. *Bird Study* 53, 92-95.
- Buchanan, G.M., Pearce-Higgins, J.W., Wotton, S.R., Grant, M.C. & Whitfield, D.P. (2003). Correlates of the change in Ring Ouzel *Turdus torquatus* abundance in Scotland from 1988-91 to 1999. *Bird Study* 50, 97-105.
- Bullock, D.J. (1985). Annual diets of hill sheep and feral goats in southern Scotland. *Journal of Applied Ecology* 22, 423-433.
- Bullock, J.M., Hill, B.C., Dale, M.P., & Silvertown, J. (1994). An Experimental-Study of the Effects of Sheep Grazing on Vegetation Change in a Species-Poor Grassland

and the Role of Seedlings Recruitment into Gaps. *Journal of Applied Ecology* 31, 493-507.

Bunnell, F.L. & Gillingham, M.P. (1985). Foraging behavior: Dynamics of dining out. In: Hudson R.J. and White R.G. (eds). *Bioenergetics of Wild Herbivores*. pp. 53-79. Boca Raton, Florida: CRC Press.

Burfield, I.J. (2002). The breeding ecology and conservation of the ring ouzel *Turdus torquatus* in Britain. PhD thesis, University of Cambridge.

Burges, S. & Weildinger, K. (2000). Estimation of calcium intake by Meadow Pipit nestlings in an acidified area. *Journal of Ornithology* 31, 426–429.

Burges, S. (1994). Segregation of diet in water pipit *Anthus spinoletta* and Meadow Pipit *Anthus pratensis* nestlings in an area damaged by air pollution. *Folia Zoologica* 43, 43–48.

Burnham, K.P., & Anderson, D.R. (2002). *Model Selection and Inference: A Practical Information-Theoretic Approach*. Fort Collins, Colorado.

Butterfield, J. & Coulson, J.C. (1975). Insect food of adult red grouse *Lagopus lagopus scoticus* Lath. *Journal of Animal Ecology* 44, 601–608.

Butterfield, J. & Coulson, J.C. (1983). The Carabid communities on peat and upland grasslands in northern England. *Holarctic Ecology* 6, 163–174.

Byrkjedal, I. (1980). Summer food of the Golden Plover *Pluvialis apricaria* at Hardangervidda, southern Norway. *Holarctic Ecology* 3, 40–58.

Calladine, J., Baines, D. & Warren, P. (2002). Effects of reduced grazing on population density and breeding success of black grouse in northern England. *Journal of Applied Ecology* 39, 772–780.

Cayford, J.T., Tyler, G. & MacIntosh-Williams, L. (1989). *The Ecology and Management of Black Grouse in Conifer Forests in Wales*. RSPB, Sandy.

Chinery, M. (1986). *A Field Guide to the Insects of Britain and Northern Europe*, 2nd edn. London: Collins.

Clarke, J.L., Welch, D. & Gordon, I.J. (1995a). The influence of vegetation pattern on the grazing of heather moorland by red deer and sheep. I. The location of animals on grass/heather mosaics. *Journal of Applied Ecology* 32, 166-176.

Clarke, J.L., Welch, D. & Gordon, I.J. (1995b). The influence of vegetation pattern on the grazing of heather moorland by red deer and sheep. II. The impact on heather. *Journal of Applied Ecology* 32, 177-186.

Clutton-Brock, T.H., Iason, G.R. & Guinness, F.E. (1987). Sexual segregation and density-related changes in habitat use in male and female red deer (*Cervus elaphus*). *Journal of Zoology* 211, 275-289.

Colquhoun, I.R. (1970). *The Grazing Ecology of Red Deer and Blackface Sheep in Perthshire, Scotland*. The Grazing Ecology of Red Deer and Blackface Sheep in Perthshire, Scotland. University of Edinburgh, Edinburgh.

Common, T.G., Grant, S.A., Armstrong, R.H. & Torvell, L. (1997). The effects of level of *Molinia* utilization on diet selection and herbage intake by cattle grazing *Molinia* grassland. *Grass and Forage Science* 52, 207-218.

Common, T.G., Wright, I.A. & Grant, S.A. (1998). The effect of grazing by cattle on animal performance and floristic composition in *Nardus*-dominated swards. *Grass and Forage Science* 53, 260-269.

Conington, J., Bishop, S.C., Waterhouse, A., & Simm, G. (2000). A bio-economic approach to estimating breeding values in UK hill sheep. *British Society of Animal Science Annual meeting*, Paper no 39.

Conington, J., Bishop, S.C., Waterhouse, A., & Simm, G. (2004). A bio-economic approach to derive economic values for pasture-based sheep genetic improvement programmes. *Journal of Animal Science* 82, 1290-1304.

Coughenour, M., Reid, R., & Thornton, P. (2000). The SAVANNA model: providing solutions for wildlife preservation and human developments in East Africa and the Western United States. <http://www.futureharvest/news/savannarelease.pdf>

Coulson, J.C. & Butterfield, J.E.L. (1985). The invertebrate communities of peat and upland grasslands in the north of England and some conservation implications. *Biological Conservation* 34, 197-225.

Coulson, J.C. & Butterfield, J.E.L. (1986). The spider communities on peat and upland grasslands in northern England. *Holarctic Ecology* 9, 229–239.

Coulson, J.C. & Whittaker, J.B. (1978). Ecology of moorland animals. In Heal, O.W. & Perkins, D.F. (eds) *Production Ecology of British Moors and Montane Grasslands*: 52–93. Berlin: Springer–Verlag.

Coulson, J.C. (1962). The biology of *Tipula subnodicornis* Zetterstedt, with comparative observations on *Tipula paludosa* Meigen. *Journal of Animal Ecology* 31, 1–21.

Coulson, J.C. (1988). The structure and importance of invertebrate communities on peatlands and moorlands, and effects of environmental and management changes. In Usher, M.B. & Thompson, D.B.A. (eds) *Ecological Change in the Uplands*: 365–380. Oxford: Blackwell Scientific Publications.

Coulson, J.C., Butterfield, J.E.L. & Henderson, E. (1990). The effect of open drainage ditches on the plant and invertebrate communities of moorland and on the decomposition of peat. *Journal of Applied Ecology* 27, 549–561.

Cramp, S. & Simmons, K.E.L. (1980). *The Birds of the Western Palearctic*, Vol. 2. Oxford: Oxford University Press.

Cramp, S. & Simmons, K.E.L. (1983). *The Birds of the Western Palearctic*, Vol. 3. Oxford: Oxford University Press.

Cramp, S. (ed.) (1988). *The Birds of the Western Palearctic*, Vol. 5. Oxford: Oxford University Press.

Crawley, M.J. (1993). *GLIM for Ecologists*. Blackwell Scientific Publications Oxford.

Crosson P., O’Kiely P., O’Mara F.P. and Wallace M. (2006) The development of a mathematical model to investigate Irish beef production systems. *Agricultural Systems* 89, 349-370.

Cummins, S. & O’Halloran, J. (2002). An assessment of the diet of nestling Stonechats *Saxicola torquata* using compositional analysis. *Bird Study* 49, 139–145.

Danilov, N.N., Ryzhanovskiy, V.N. & Ryabitsev, V.K. (1984). *Ptitsy Yamala*. Moscow: Nauka.

Davies, O.D., Wildig, J., Jones, D.L., (2000). The implications of Environmentally Sensitive Area prescriptions on livestock performance off rough grazings. In: *Grazing Management: the Principles and Practice of Grazing, for Profit and Environmental Gain, within Temperate Grassland Systems*, A.J. Rook & P.D. Penning (eds). BGS Occasional Symposium No. 34, 153-154.

Defra (2004). An assessment of the impacts of hill farming in England on the economic, environmental and social sustainability of the uplands and more widely. Defra, UK.

Defra (2005). UK crop areas and livestock numbers.

Defra (2006). Agricultural Change and Environmental Observatory Programme Annual Review. www.defra.gov.uk/farm/policy/observatory/annualreview.htm

Demment M.W. & Laca E.A. (1993). The grazing ruminant: models and experimental techniques to relate sward structure and intake. *Proceedings of the World Conference on Animal Production*, Edmonton, Canada, 1993. pp. 439-460.

Demment, M.W. & Van Soest, P.J. (1985). A nutritional explanation for body-size patterns of ruminant and non-ruminant herbivores. *American Naturalist* 125, 641-675.

Dennis, P., Elston, D., Evans, D.M., Evans, S.A., Gordon, I.J., Grant, M., Kunaver, A., Marquiss, M., Mayes, R., McCracken, D.I., Pakeman, R., Pearce-Higgins, J., Redpath, S.M., Skartveit, J., Stephen, L., Benton, T. & Bryant, D. (2005). *Effects of Grazing Management on Upland Bird Populations: Disentangling Habitat Structure and Arthropod Food Supply at Appropriate Spatial Scales (GRUB)*. Unpublished report to The Scottish Executive Environment and Rural Affairs Department, Edinburgh.

Dennis, P., Young, M.R. & Bentley, C. (2001). The effects of varied grazing management on epigeal spiders, harvestmen and pseudoscorpions of *Nardus stricta* grassland in upland Scotland. *Agriculture Ecosystems and Environment* 86, 39–57.

Dennis, P., Young, M.R. & Gordon, I.J. (1998). Distribution and abundance of small insects and arachnids in relation to structural heterogeneity of grazed, indigenous grasslands. *Ecological Entomology* 23, 253–264.

Dennis, P., Young, M.R., Howard, C.L. & Gordon, I.J. (1997). The response of epigeal beetles (Col: Carabidae, Staphylinidae) to varied grazing regimes on upland *Nardus stricta* grasslands. *Journal of Applied Ecology* 34: 433–443.

Dodgshon, R.A. & Olsson, G.A. (2006). Heather moorland in the Scottish Highlands: the history of a cultural landscape, 1600–1880. *Journal of Historical Geography* 32, 21-37.

Donald, P.F., Muirhead, L.B., Buckingham, D.L., Evans, A.D., Kirby, W.B. & Gruar, D.J. (2001). Body condition, growth rates and diet of Sky Lark *Alauda arvensis* nestlings on lowland farmland. *Ibis* 143, 658–669.

Donnelly, J.R., Moore, A.D. & Freer, M. (1997). GRAZPLAN: Decision support systems for Australian grazing enterprises. 1. Overview of the GRAZPLAN project, and a description of the MetAccess and LambAlive DSS. *Agricultural Systems* 54, 57-76.

Dorzhev, T.Z. & Khertuev, V.N. (1992). *Ekologiya kamenok v Zabaikalie*. Ulan-Ude. Buryatian Scientific Centre of Siberian Branch of Russia Academy of Science 148.

Downie, I.S., Butterfield, J.E.L. & Coulson, J.C. (1995). Habitat preferences of sub-montane spiders in northern England. *Ecography* 18, 51–61.

Dunne, F. & Doyle, G. (1998). Changes in *Molinia*-dominated vegetation due to cattle grazing in Killarney, Co. Kerry. *Biology and Environment* 98B, 45.

El Aich, A. & Waterhouse, A. (1999). Small ruminants in environmental conservation. *Small Ruminant Research* 34, 271-287.

English Nature (2001). *State of nature: the upland challenge*. English Nature, Peterborough.

Equihua, M. (1990). Fuzzy clustering of ecological data. *Journal of Ecology* 78, 519-534.

Evans, D.M., Redpath, S., Evans, S.A., Gardner, C.J., Elston, D.A., Dennis, P. & Pakeman, R.J. (2006). Low intensity, mixed livestock grazing improves the breeding abundance of a common insectivorous passerine. *Biology Letters* 2, 636-638.

Felton, M., & Marsden, J.H. (1990). *Heather regeneration in England and Wales*. Nature Conservancy Council, Peterborough.

Fenton, E.W. (1935). The Transition from woodland and moorland to grassland in the Spey Valley and elsewhere. *Journal of Ecology* 23, 56-68.

Fielding, A. H. & Haworth, P. F. (1995). Testing the generality of bird-habitat models. *Conservation Biology* 9, 1466-1481.

Filimonova, N.P. (1972). Materialy o pitanii tetereva i seroi kuropatki v Kustanaiskoi oblasti. Novosibirsk. Moscow: Nauka.

Finlayson J.D., Cacho O.J. & Bywater A.C. (1995). A simulation model of grazing sheep. 1. Animal growth and intake. *Agricultural Systems* 48, 1-25.

Flores, E.R., Provenza, F.D. & Balph, D.F. (1989). The effect of experience on the foraging skill of lambs: Importance of plant form. *Applied Animal Behaviour Science* 23, 285-291.

Focardi, S. & Marcellini, P. (1995). A mathematical framework for optimal foraging of herbivores. *Journal of Mathematical Biology* 33, 365-387.

France, J., Dhanoa, M.S., Theodorou, M.K., Lister, S.J., Davies, D.R. & Isac, D. (1993). A model to interpret gas accumulation profiles associated with in vitro degradation of ruminal feeds. *Journal of Theoretical Biology* 163, 99-111.

Fraser, M.D. & Gordon, I.J. (1997). The diet of goats, red deer and South American camelids feeding on three contrasting Scottish upland vegetation communities. *Journal of Applied Ecology* 34, 668-686.

Fraser, M.D., Fychan, R. & Jones, R. (2005). The effect of harvest date and inoculation on the yield and fermentation characteristics of two varieties of white lupin (*Lupinus albus*) when ensiled as a whole-crop. *Animal Feed Science and Technology* 120, 203-216.

Fraser, M.D., Theobald, V.J. & Moorby, J.M. (2006). Determining the diet composition of animals grazing complex swards using n-alkanes and long-chain fatty alcohols. *Ecological Applications* 16, 1901-1910.

Fraser, M.D., Vale, J.E., Evans, J.G., Theobald, V.J. & Davies, D.W.R. (2007). Performance of Limousin-cross and Belted Galloway suckler cows and calves when grazing improved upland swards and Molinia-dominant semi-natural rough grazing. In: Hopkins, J.J. (ed.) *High Value Grassland: Providing Biodiversity, a Clean*

Environment and Premium Products. BGS Symposium 38, 147-151. British Grassland Society, Cirencester.

Freer, M., Moore, A.D. & Donnelly, J.R. (1997). GRAZPLAN: Decision support systems for Australian grazing enterprises. 2. The animal biology model for feed intake, production and reproduction, and the GrazFeed DSS. *Agricultural Systems* 54, 77-126.

Fuller, R.J. & Gough, S.J. (1999). Changes in sheep numbers in Britain: implications for bird populations. *Biological Conservation* 91, 73-89.

Galbraith, H. (1989). The diet of Lapwing *Vanellus vanellus* chicks on Scottish farmland. *Ibis* 131, 80-84.

Gardner, S.M. & Usher, M.B. (1989). Insect abundance on burned and cut upland *Calluna* heath. *Entomologist* 108, 145-200.

Gardner, S.M. (1991). Ground beetle (Coleoptera: Carabidae) communities on upland heath and their association with heathland flora. *Journal of Biogeography* 18, 281-289.

Gardner, S.M. (2002) Managing upland vegetation for sheep and conservation. In: *Conservation pays? Reconciling environmental benefits with profitable grassland systems* (ed. J. Frame). Occasional Symposium of the British Grassland Society, No. 36, pp. 115-118. British Grassland Society, Reading.

Gardner, S.M., Hartley, S.E., Davies, A. & Palmer, S.C.F. (1997). Carabid communities on heather moorlands in northeast Scotland: the consequences of grazing pressure for community diversity. *Biological Conservation* 81, 275-286.

Gardner, S.M., Hetherington, S.L. & Allen, D. (2002). Assessment of vegetation change and *Calluna/Nardus* interactions in relation to spatial variation in grazing pressure on upland moor. Unpublished Final Report to DEFRA (Contract BD1211).

Gardner, S.M., Ross, S.Y. & Adamson (2001). Assessment of the effect of spatial variation in sheep grazing on *Calluna vulgaris-Molinia caerulea* interactions. Unpublished Final Report to DEFRA (Contract BD1218).

Garnett, M.C., MacFadzean, S. & Townsend, M. (1988). The Foraging Ecology of the Whinchat *Saxicola rubetra* in Wales. Unpublished RSPB report. RSPB, Sandy.

Gibbons, D.W., Avery, M.I., Baillie, S.R., Gregory, R.D., Kirby, J., Porter, R.F., Tucker, G.M. & Williams, G. (1996). Bird species of Conservation Concern in the United Kingdom, Channel Islands and Isle of Man: revising the Red Data list. *RSPB Conservation Review* 10, 7-18.

Gibbons, D.W., Reid, J.B. & Chapman, R.A. (1993). *The New Atlas of Breeding Birds in Britain and Ireland: 1988-1991*. Poyser, London.

Gibson, L.A., Wilson, B.A., Cahill, D.M. & Hill, J. (2004). Spatial prediction of rufous bristlebird habitat in a coastal heathland: a GIS-based approach *Journal of Applied Ecology* 41, 213–223.

Gilbert, O.L. & Anderson, P. (1998). *Habitat creation and repair* Oxford University Press, Oxford.

Gimingham, C.H. (1985). Age-related interactions between *Calluna vulgaris* and phytophagous insects. *Oikos* 44, 12–16.

Gimingham, C.H. (1989). Heather and Heathlands. *Botanical Journal of the Linnean Society* 101, 263-268.

Gimingham, C.H. (1998). Introduction: Ecological change in the uplands. In: Usher, M.B., Thompson, D.B.A. (eds) *Ecological Change in the Uplands*, Blackwell Scientific Publications, Oxford.

Gordon, I.J. & Hutchings, N.J. (1993). The development of sustainable ruminant livestock grazing systems: the role of modelling. *Proceedings of the World Conference on Animal Production, Edmonton, Canada, 1993*. pp. 413-438.

Gordon, I.J. & Illius, A.W. (1988). Incisor arcade structure and diet selection in ruminants. *Functional Ecology* 2, 15-22.

Gordon, I.J., Hester, A.J. & Festa-Bianchet, M. (2004). The management of wild large herbivores to meet economic, conservation and environmental objectives. *Journal of Applied Ecology* 41, 1020-1031.

Gosler, A. (1993). *The Great Tit*. London: Hamlyn.

Gower, J.C. (1971). Statistical methods of comparing different multivariate analyses. In: Hodson, F.R. et al. (Eds.) *Mathematics in the Archaeological and Historical Sciences*, Edinburgh University Press.

Graf, R.F., Bollmann, K., Sachot, S., Suter, W. & Bugmann, W. (2006). On the generality of habitat distribution models: a case study of capercaillie in three Swiss regions. *Ecography* 29, 319–328.

Grant, M.C., Orsman, C., Easton, J., Lodge, C., Smith, M., Thompson, G., Rodwell, S. & Moore, N. (1999). Breeding success and causes of breeding failure of curlew *Numenius arquata* in Northern Ireland. *Journal of Applied Ecology* 36, 59-74.

Grant, S.A. & Campbell, D.R. (1978). Seasonal variation in in vitro digestibility and structural carbohydrate content of some commonly grazed plants of blanket bog. *Journal of the British Grassland Society* 33, 167-173.

Grant, S.A. & Hodgson, J. (1986). Grazing effects on species balance and herbage production in indigenous plant communities. In: Gudmundsson O. (ed). *Grazing Research at Northern Latitudes*. pp. 69-77. London: Plenum Publishing Corporation.

Grant, S.A. & Maxwell, T.J. (1988). Hill vegetation and grazing animals: the biology and definition of management options. In: Usher M.B. and Thompson D.B.A. (eds). *Ecological Change in the Uplands*. pp. 201-214. Oxford: Blackwell Scientific Publications.

Grant, S.A., & Hunter, R.F. (1968). Interactions of grazing and burning on heather moors and their implications in heather management. *Journal of the British Grassland Society* 23, 285-293.

Grant, S.A., Barthram, G.T., Lamb, W.I.C. & Milne, J.A. (1978). Effects of season and level of grazing on the utilisation of heather by sheep. I. Responses of the sward. *Grass and Forage Science* 33, 289-300.

Grant, S.A., Bolton, G.R. & Torvell, L. (1985a). The responses of blanket bog vegetation to controlled grazing by hill sheep. *Journal of Applied Ecology* 22, 739-751.

Grant, S.A., Hamilton, W.J. & Souter, C. (1981). The responses of heather dominated vegetation in North East Scotland to grazing by red deer. *Journal of Ecology* 69, 189-204.

Grant, S.A., Hunter, R.F. & Cross, C. (1963). The effects of burning *Molinia*-dominant communities. *Grass and Forage Science* 18, 249-257.

Grant, S.A., Lamb, W.I.C., Kerr, C.D. & Bolton, G.R. (1976). The utilisation of blanket bog vegetation by grazing sheep. *Journal of Ecology* 13, 857-869.

Grant, S.A., Suckling, D.E., Smith, H.K., Torvell, L., Forbes, T.D.A., & Hodgson, J. (1985b). Comparative studies of diet selection by sheep and cattle: the hill grasslands. *Journal of Ecology* 73, 987-1004.

Grant, S.A., Suckling, S.A., Smith, H.K., Torvell, L., Forbes, T.D.A. & Hodgson, J. (1987). Comparative studies of diet selection by sheep and cattle: blanket bog and heather moor. *Journal of Ecology* 75, 947-960.

Grant, S.A., Torvell, L., Common, T.G., Sim, E.M. & Small, J.L. (1996a). Controlled grazing studies on *Molinia* grassland: effects of different seasonal patterns and levels of defoliation on *Molinia* growth and responses of swards to controlled grazing by cattle. *Journal of Applied Ecology* 33, 1267-1280.

Grant, S.A., Torvell, L., Sim, E.M., Small, J.L. & Armstrong, R.H. (1996b). Controlled grazing studies on *Nardus* grassland: effects of between-tussock sward height and species of grazer on *Nardus* utilisation and floristic composition in two fields in Scotland. *Journal of Applied Ecology* 33, 1053-1064.

Grant, S.A., Torvell, L., Sim, E.M., Small, J.L. & Elston, D.A. (1996c). Seasonal pattern of leaf growth and senescence of *Nardus stricta* and responses of tussocks to differing severity, timing and frequency of defoliation. *Journal of Applied Ecology* 33, 1145-1155.

Green, R.E. (1986). Breeding Waders of the Somerset Moors: Factors Affecting Their Distribution and Breeding Success. Unpublished report to NCC. NCC, Peterborough.

Green, R.E. (2006). Factors Affecting the Population Density of the Corncrake *Crex crex* in Britain and Ireland. *Journal of Applied Ecology* 33, 237-248.

Green, R.E., Hirons, G.J.M. & Cresswell, B.H. (1990). Foraging habitats of female common snipe *Gallinago gallinago* during the incubation period. *Journal of Applied Ecology* 27, 325-335.

Gregory, R.D., Wilkinson, N.I., Noble, D.G., Robinson, J.A., Brown, A.F., Hughes, J., Proctor, D., Gibbons, D.W. & Galbraith, C.A. (2002). The population status of birds in the United Kingdom, Channel Islands and Isle of Man: an analysis of conservation concern 2002-07. *British Birds* 95, 410-448.

- Greig-Smith, P.W. & Quicke, D.L.J. (1983). The diet of nesting Stonechats. *Bird Study* 30, 47–50.
- Griffiths, J.B., Henman, D., Jones, D.L., Wildig, J., Rushton, S.P. & Sanderson, R. (1997). Stocking rate comparisons on semi-natural rough grazings in the Cambrian Mountains. In *Grassland Management in Environmentally Sensitive Areas* (Ed. R.D. Sheldrick) BGS Occasional Symposium No. 32: 232-233.
- Grime, J.P., Hodgson, J.G. & Hunt, R. (1988). *Comparative Plant Ecology: a functional approach to common British species*. Unwin Hyman, London.
- Grime, J.P., Hodgson, J.G., & Hunt, R. (1996). *Comparative Plant Ecology. A functional approach to common British species*. The Ipswich Book Company Ltd, Ipswich, Suffolk.
- Gubin, B.M. & Kovshar, A.F. (1985). K biologii obyknovnoy kamenki v vysokogor'ie Tien-Shanya. *Ornitologiya* 20, 53–59.
- Hadjigeorgiou, I., Osoro, K., Fragoso de Almeida, J.P. & Molle, G. (2005). Southern European grazing lands: Production, environmental and landscape management aspects. *Livestock Production Science* 96, 51-59.
- Hagvar, S. & Ostbye, E. (1976). Food habits of the Meadow Pipit *Anthus pratensis* in alpine habitats at Hardangervidda, south Norway. *Norwegian Journal of Zoology* 24, 53–67.
- Haines-Young, R.H. et. al. (2000). *Accounting for nature: assessing habitats in the UK countryside*. Department of the Environment, Transport, and the Regions. London.
- Hancock, M., Baines, D., Gibbons, D., Etheridge, B. & Shepherd, M. (1999). Status of male black grouse *Tetrao tetrix* in Britain in 1995-96. *Bird Study* 46, 1-15.
- Hartley, S.E. & Amos, L. (1999). Competitive interactions between *Nardus stricta* L. and *Calluna vulgaris* (L.) Hull: the effect of fertilizer and defoliation on above- and below-ground performance. *Journal of Ecology* 87, 330-340.
- Hartley, S.E. & Mitchell, R.J. (2005). Manipulation of nutrients and grazing levels on heather moorland: changes in *Calluna* dominance and consequences for community composition. *Journal of Ecology* 93, 990-1004.

Hartley, S.E., (1997). The effects of grazing and nutrient inputs on grass-heather competition. *Botanical Journal of Scotland* 49, No.2, 315-324.

Hartley, S.E., Gardner, S.M. & Mitchell, R.J. (2003). Indirect effects of grazing and nutrient addition on the hemipteran community of heather moorlands. *Journal of Applied Ecology* 40, 793–803.

Haworth, P.F. & Thompson, D.B.A. (1990). Factors associated with the breeding distribution of upland birds in the south Pennines. England. *Journal of Applied Ecology* 27, 562-577.

Hayes, G.F. & Holl, K.D. (2003). Cattle grazing impacts on annual forbs and vegetation composition of mesic grasslands in California. *Conservation Biology* 17, 1694-1702.

Haysom, K.A. & Coulson, J.C. (1998). The Lepidoptera fauna associated with *Calluna vulgaris*: effects of plant architecture on abundance and diversity. *Ecological Entomology* 23, 377–385.

Henderson, D.C. (1995). Condition score targets for mating. The veterinary book for sheep farmers. Farming Press. ISBN 0-85236-189-0.

Hester, A.J. & Baillie, G.J. (1998). Spatial and temporal patterns of heather use by sheep and red deer within natural heather/grass mosaics. *Journal of Applied Ecology* 35, 772-784.

Hester, A.J. & Sydes, C. (1992). Changes in burning of Scottish heather moorland since the 1940s from aerial photographs. *Biological Conservation* 60, 25–30.

Hester, A.J., Gordon, I.J., Baillie, G.J. & Tappin, E. (1999). Foraging behaviour of sheep and red deer within natural heather grass mosaics. *Journal of Applied Ecology* 36, 133-146.

Hester, A.J., Mitchell, F.J.G., Gordon, I.J. & Baillie, G.J. (1996). Activity patterns and resource use by sheep and red deer grazing across a grass/heather boundary. *Journal of Zoology* 240, 609-620.

Hetherington, S.L. (2000). The use of self-help feed blocks as an aid to grazing and vegetation management of semi-natural rough grazing. *Aspects of Applied Biology* 58, 137-142.

Hetherington, S.L., McLean, B.M.L., Gardner, S.M., Wildig, J. & Griffiths, J.B. (2002). The impact of Environmentally Sensitive Areas policy in relation to conservation and farming objectives. In: Conservation pays? Reconciling environmental benefits with profitable grassland systems (ed. J. Frame). Occasional Symposium of the British Grassland Society, No. 36, pp. pp. 85-88. British Grassland Society, Reading.

Hewson, R. (1976). Grazing by mountain hares *Lepus timidus* L., red deer *Cervus elaphus* L. and red grouse *Lagopus L. scoticus* on heather moorland in north-east Scotland. *Journal of Applied Ecology* 13, 657-666.

Hill, M.O. & Gauch, H.G. (1980). Detrended correspondence analysis, an improved ordination technique. *Vegetatio* 42, 47-58.

Hill, M.O., Evans, D.F., & Bell, S.A. (1992). Long-term effects of excluding sheep from hill pastures in North Wales. *Journal of Ecology* 80, 1-13.

Hobbs, R.J. & Gimingham, C.H. (1987). Vegetation, fire and herbivore interactions in Scottish heathland communities. II. Post-fire vegetation development. *Journal of Ecology* 72, 585-610.

Hobson, P.N. & Wallace, R.J. (1982). Microbial ecology and activities in the rumen. *Critical Reviews in Microbiology* 9, 165 - 225.

Hodgson, J. & Eadie, J. (1986). Vegetation resources and animal nutrition in hill areas: agricultural and environmental implications. In: O'Toole M. (ed). Hill Farming Symposium, Galway 1984. pp. 118-133. Dublin: An Foras Taluntais.

Hodgson, J. (1982). Influence of sward characteristics on diet selection and herbage intake by the grazing animal. In: Hacker J.B. (ed). Nutritional Limits to Animal Production from Pastures.

Hodgson, J., Forbes, T.D.A., Armstrong, R.H., Beattie, M.M. & Hunter, E.A. (1991). Comparative Studies of the Ingestive Behaviour and Herbage Intake of Sheep and Cattle Grazing Indigenous Hill Plant Communities. *Journal of Applied Ecology* 28, 205-227.

Hofmann, R.R. (1989). Evolutionary steps of ecophysiological adaptation and diversification of ruminants: a comparative view of their digestive system. *Oecologia* 78, 443-457.

Holmes, R.T. (1966). Feeding ecology of the red backed sandpiper *Calidris alpina* in arctic Alaska. *Ecology* 47, 32–45.

Holmes, R.T. (1970). Differences in population density, territoriality, and food supply of dunlin on arctic and subarctic tundra. In Watson, A. (ed.) *Animal Populations in Relation to Their Food Resources*: 303–319. Symposium of the British Ecological Society. Oxford: Blackwell.

Hoodless, A.N., Inglis, J.G., & Baines, D. (2006). Effects of weather and timing on counts of breeding Snipe *Gallinago gallinago*. *Bird Study* 53, 205-212.

Hope, D., Picozzi, N., Catt, D.C. & Moss, R. (1996). Effects of reducing sheep grazing in the Scottish Highlands. *Journal of Range Management* 49, 301-310.

<http://statistics.defra.gov.uk/esg/publications/auk/2004/excel.asp>

Hudson, P.J. (1992). *Grouse in Space and Time*. Game Conservancy Trust, Fordingbridge.

Hulme, P.D., Merrell, B.G., Torvell, L., Fisher, J.M., Small, J.L. & Pakeman, R.J. (2002). Rehabilitation of degraded *Calluna vulgaris* (L.) Hull-dominated wet heath by controlled sheep grazing. *Biological Conservation* 107, 351-363.

Hulme, P.D., Pakeman, R.J., Torvell, L., Fisher, J.M. & Gordon, I.J. (1999). The effects of controlled sheep grazing on the dynamics of upland *Agrostis-Festuca* grassland. *Journal of Applied Ecology* 36, 886-900.

Hunter, R.F. (1962). Hill sheep and their pasture: a study of sheep-grazing in south east Scotland. *Journal of Ecology* 50, 651-680.

Hutchings, N.J. (1991). A model of a grass sward continuously grazed to a constant mean height. *Ecological modelling* 59, 73-91.

Hyslop, J.J. & Merrell, B.G. (2002). Hill sheep physical and financial performance when managed according to environmentally sensitive area (ESA) prescriptions. In: *Proceedings of the SAC Workshop: Biodiversity, plant structure and vegetation heterogeneity: interactions with the grazing herbivore*. Published on CD-ROM by SAC, Hill and Mountain Research Centre, Food Systems Division, Kirkton Farm, Crianlarich, Perthshire FK20 8RU.

Illius, A.W. & Gordon, I.J. (1987). The allometry of food intake in grazing ruminants. *Journal of Animal Ecology* 56, 989-999.

Illius, A.W. & Gordon, I.J. (1991). Prediction of intake and digestion in ruminants by a model of rumen kinetics integrating animal size and plant characteristics. *Journal of Agricultural Science* 116, 145-157.

Illius, A.W. (1986). Foraging behaviour and diet selection. In: Gudmundsson O. (ed) *Grazing Research at Northern Latitudes. Proceedings of a NATO Advanced Research Workshop, Hvanneyri, 1985.* pp. 227-236. New York: Plenum Press.

Illius, A.W. (1989). Allometry of food intake and grazing behaviour with body size in cattle. *Journal of Agricultural Science* 113, 259-266.

Jacobs, J. (1974). Quantitative measurement of food selection. A modification of the forage ratio and Ivlev's electivity index. *Oecologia* 14, 413-417.

Jalal, M.A.F., Read, D.J., & Haslam, E. (1982). Phenolic composition and its seasonal variation in *Calluna vulgaris*. *Phytochemistry* 21, 1397-1401.

Janecek, S. & Leps, J. (2005). Effect of litter, leaf cover and cover of basal internodes of the dominant species *Molinia caerulea* on seedling recruitment and established vegetation. *Acta Oecologica* 28, 141-147.

Janssen S. and van Ittersum M.K. (2007) Assessing farm innovations and responses to policies: a review of bio-economic farm models. *Agricultural Systems* 94, 622-636.

Jarman, P.J. & Sinclair, A.R.E. (1979). Feeding strategy and the pattern of resource partitioning in ungulates. In: Sinclair A.R.E. and Norton-Griffiths M. (eds). *Serengeti, Dynamics of an Ecosystem.* pp. 130-163. Chicago: University of Chicago Press.

Jarman, P.J. (1974). The social organisation of antelope in relation to their ecology. *Behaviour* 48, 215-267.

Jenny, M. (1990). Nahrungsoekologie der Feldlerche *Alauda arvensis* in einer intensiv genutzten Agrarlandschaft des schweizerischen Mittellandes. *Der Ornithologische Beobachter* 87, 31-53.

Jerram, E. (2000). Exmoor Heaths. Survey of National Vegetation Classification Communities. Unpublished report for English Nature.

JNCC (2004). Upland Habitat Common Standard Monitoring Guidance. JNCC, Peterborough.

Johnson, I.R. & Parsons, A.J. (1985). Use of a model to analyse the effects of continuous grazing management on seasonal patterns of grass production. *Grass and Forage Science* 40, 449-458.

Kapoor-vijay, P. & Usher, M.B. (1993). Identification of Key Species for Conservation and Socio- Economic Development. Commonwealth Secretariat, London.

Kastdalen, L. & Wegge, P. (1985). Animal food in Capercaillie and Black Grouse chicks in south east Norway – a preliminary report. In Lovel, T.W.I. & Hudson, P.J. (eds) *Proceedings of the Third International Grouse Symposium*: 499–513. Reading, UK: World Pheasant Association.

Kirkham, F.W. & Milne, J.A. (2000). Progress towards defining ecologically-sustainable grazing management: the 'Moorland Biomass' and 'Heather Suppression' projects. *Aspects of Applied Biology* 58, 151-158.

Kleijn, D. & Sutherland, W. J. (2003). How effective are European agri-environment schemes in conserving and promoting biodiversity? *Journal of Applied Ecology* 40, 947-969.

Kluyver, H.N. & Tinbergen, L. (1953). Territory and the regulation of density in titmice. *Archives Néerlandaises de Zoologie* 10, 265–289.

Kondrat'yev, A.Y. & Kondrat'yeva, L.F. (1988). Ispol'zovanie rezul'tatov gelmintologicheskikh vskrytiy dlya analiza pishchevykh svyazei kulikov. Moscow: Nauka.

Kostin, Y.V. (1983). Ptitsy Kryma. Moscow: Nauka.

Kumari, A. (1958). Rabakurvitsaliste toitumisest. – Rmt-s: E. Kumari toim. Ornitoloogiline kogumik, 1. Eesti NSV Teaduste Akadeemia Toimetuskirjastusnõukogu, Tartu. Lk., 195–215.

Laiolo, P., Dondero, F., Ciliento, E. & Rolando, A. (2004). Consequences of pastoral abandonment for the structure and diversity of alpine avifauna. *Journal of Applied Ecology* 41, 294-304.

Laurenroth, W.K., Hunt, H.W., Swift, D.M. & Singh, J.S. (1986). Estimating aboveground net primary production in grasslands: a simulation approach. *Ecological Modelling* 33, 297-314.

Lehtonen H., Aakkula J., and Rikkonen P. (2005) Alternative agricultural policy scenarios, sector modeling and indicators: A sustainability assessment. *Journal of Sustainable Agriculture* 26(4): 63-93.

Leps, J. & Smilauer, P. (2003). *Multivariate analysis of ecological data using CANOCO*. Cambridge University Press.

Lima, S.L. & Dill, L.M. (1990). Behavioral decisions made under the risk of predation: A review and prospectus. *Canadian Journal Zoology* 68, 619-640.

Linsley, M.D. (1999). *The Effects of Intensive Agriculture on the Breeding Ecology of the Lapwing Vanellus vanellus*. PhD thesis, Liverpool John Moores University.

Littlewood, N.A., Pakeman, R.J. & Woodin, S.J. (2006). A field assessment of the success of moorland restoration in the rehabilitation of whole plant assemblages. *Applied Vegetation Science* 9, 295-306.

Lovegrove, R., Shrubbs, M. & Williams, I. (1995). *Silent Fields: the Current Status of Farmland Birds in Wales*. RSPB, Newtown.

Lowman, B.G., Hinks, C.E., Hunter, E.A. & Scott, N.A. (1996). Effect of breed type, sex, method of rearing and winter nutrition on lifetime performance and carcass composition in a 20-month beef system: Grazing performance. *Animal Science* 63, 215-222.

Lowman, B.G., Scott, N.A. & Somerville S.H. (1976). Condition scoring of cattle. Rev. Ed. Bulletin, East of Scotland College of Agriculture, No.6.

Macdonald, A. J., Kirkpatrick, A.H., Hester, A.J. & Sydes, C. (1995). Regeneration by natural layering of heather (*Calluna vulgaris*): frequency and characteristics in upland Britain. *Journal of Applied Ecology* 32, 85-99.

Mackey, E.C., Shewry, M.C. & Tudor, G.J. (1998). *Land Cover Change: Scotland from the 1940s to the 1980s*. HMSO, Edinburgh.

MAFF (1994). Body Condition Scoring of Sheep (leaflet PB1875). MAFF Publications, London.

Mal'chevskiy, A.S. & Pukinskiy, Y.B. (1983). Ptitsy Leningradskoi oblasti i sopredel'nykh territoriy 2. Leningrad: Leningrad University Press.

Mandaluniz, N., Oregui, L.M. & Aldezabal, A. (2005). Utilisation of gorse and heather communities by cattle in mountain grazing. In: Sylvopastoralism and Sustainable Land Management (eds Mosquera-Losada, M. R.; Rigueiro-Rodriguez, A.; McAdam, J.), pp.157-159. CABI Publishing, Wallingford.

Marrs, R.H. & Welch, D. (1991). Moorland wilderness: the potential effects of a reduced grazing pressure Report to Department of the Environment, London, UK.

Marrs, R.H., Phillips, J.D.P., Todd, P.A., Ghorbani, J., & Le Duc, M.G. (2004). Control of *Molinia caerulea* on upland moors. *Journal of Applied Ecology* 41, 398-411.

Martin, L.M. & Wilsey, B.J. (2006). Assessing grassland restoration success: relative roles of seed addition and native ungulate activities. *Journal of Applied Ecology* 43, 1098-1109.

Matthews K.B., Wright I.A., Buchan K., Davies, D.A. and Schwarz G. (2006) Assessing the options for upland livestock systems under CAP reform: Developing and applying a livestock systems model within whole-farm systems analysis. *Agricultural Systems*, **90**, 32-61.

Maxwell, T.J., Grant, S.A., Milne, J.A. & Sibbald, A.R. (1986). Systems of sheep production on heather moorland. Proceedings of a Hill Land Symposium, Galway, 1984, 187-211.

Mayes, R.W., Lamb, C.S. & Colgrove, P.M. (1986). The use of dosed and herbage n-alkanes as markers for the determination of herbage intake. *Journal of Agricultural Science* 107, 161-170.

McArthur, C., Robbins, C.T., Hagerman, A.E. & Hanley, T.A. (1993). Diet selection by a ruminant generalist browser in relation to plant chemistry. *Canadian Journal of Zoology* 71, 2236-2243.

McCall, D.G., Marshall, P.R. & Johns, K.L. (1991). An introduction to STOCKPOL: a decision support model for livestock farms. Proceedings of the International

Conference on Decision Support Systems for Rural Resource Management, Texas A & M University, April 1991. pp. 27-30.

McCracken, D.I. & Foster, G.N. (1993). Surface-active invertebrate communities and the availability of potential food for the Chough, *Pyrrhocorax pyrrhocorax* L., on pastures in northwest Islay. *Pedobiologia* 37, 141–158.

McFerran, D.M., Montgomery, W.I. & McAdam, J.H. (1994). Effects of grazing intensity on heathland vegetation and ground beetle assemblages of the uplands of County Antrim, northeast Ireland. *Biology and Environment* 94B, 41–52.

McFerran, D.M., Montgomery, W.I. & McAdam, J.H. (1994). The impact of grazing on communities of ground-dwelling spiders Araneae in upland vegetation types. *Biology and Environment* 94B, 119–126.

Merrell, B.G., Wildig, J. & Davies, O.D. (2001). Moorland conservation and hill sheep. Integrated Upland Management for Wildlife, Field Sports, Agriculture & Public Enjoyment (Phillips, J.D.P., Thompson, D.B.A. & Gruellich, W.H. – Eds), Scottish Natural Heritage, Battleby 2001, 75 – 93.

Miller, G.R. & Cummins, R.P. (2003). Soil seed banks of woodland, heathland, grassland, mire and montane communities, Cairngorm Mountains, Scotland. *Plant Ecology* 168, 255-266.

Milligan, A.L., Putwain, P.D., & Marrs, R.H. (1999). A laboratory assessment of the relative susceptibility of *Molinia caerulea* (L.) Moench and *Calluna vulgaris* (L.) Hull to a range of herbicides. *Annals of Applied Biology*, 135, 503-508.

Milligan, A.L., Putwain, P.D., Cox, E.S., Ghorbani, J., Le Duc, M.G., & Marrs, R.H. (2004). Developing an integrated land management strategy for the restoration of moorland vegetation on *Molinia caerulea*-dominated vegetation for conservation purposes in upland Britain. *Biological Conservation* 119, 371-385.

Milne, J.A. & Bagley, L. (1976). The nutritive value of diets containing different proportions of grass and heather (*Calluna vulgaris* L. Hull) to sheep. *Journal of Agricultural Science Cambridge*, 87, 599-604.

Milne, J.A. & Hartley, S.E. (2001). Upland plant communities - sensitivity to change. *Catena* 42, 333-343.

Milne, J.A. & Sibbald, A.R. (1998). Modelling of grazing systems at the farm level. *Annales de Zootechnie* 47, 407-417.

Milne, J.A. (1974). The effects of season and age of stand on the nutritive value of heather (*Calluna vulgaris*, L. Hull) to sheep. *Journal of Agricultural Science Cambridge*, 83, 281-288.

Milne, J.A. (1980). Comparative digestive physiology and metabolism of the red deer and the sheep. *Proceedings of the New Zealand Society of Animal Production* 40, 151-157.

Milne, J.A. (1998). Appendix 11 – Computer-based decision support tools to aid grazing management. In: Stuart F.E. and Eno S.G. *Grazing Management Planning for Upland Natura 2000 Sites: A Practical Manual*. pp. 127-128. Edinburgh: The National Trust for Scotland.

Milne, J.A., Bagley, L. & Grant, S.A. (1979). Effects of season and level of grazing on the utilization of heather by sheep. 2. Diet selection and intake. *Grass and Forage Science* 34, 45-53.

Milne, J.A., Macrae, J.C., Spence, A. & Wilson, S. (1978). A comparison of the voluntary intake and digestion of forages at different times of the year by the sheep and red deer (*Cervus elaphus*). *British Journal of Nutrition* 40, 347-357.

Milne, J.A., Pakeman, R.J., Kirkham, F.W., Jones, I.P. & Hossell, J.E. (2002). Biomass production of upland vegetation types in England and Wales. *Grass and Forage Science* 57, 373-388.

MLC (1983). *Getting ewes ready for tugging*. Milton Keynes, UK: Meat and Livestock Commission.

Mohamed, B.F. & Gimingham, C.H. (1970). The Morphology of Vegetative Regeneration in *Calluna vulgaris*. *New Phytologist* 69, 743-750.

Moore, A.D., Donnelly, J.R. & Freer, M. (1997). GRAZPLAN: Decision support systems for Australian grazing enterprises. 3. Pasture growth and soil moisture sub-models, and the GrassGro DSS. *Agricultural Systems* 55, 535-582.

Moreby, S.J. & Stoate, C. (2000). A quantitative comparison of neck collar and faecal analysis to determine passerine nestling diet. *Bird Study* 47, 320–331.

Morris, A.J., Whittingham, M.J., Bradbury, R.B., Wilson, J.D., Kyrkos, A., Buckingham, D.L., Evans, A.D. (2001). Foraging habitat selection by yellowhammers (*Emberiza citrinella*) nesting in agriculturally contrasting regions in lowland England. *Biological Conservation* 101, 197-210.

Moss, R. (1989). Management of heather for game and livestock. *Botanical Journal of the Linnean Society* 101, 301-306.

Murray, M.G., Lewis, A.R. & Coetzee, A.M. (1981). An evaluation of capture techniques for research on impala populations. *South African Journal of Wildlife Research* 11, 105-109.

Nechaev, A.V. (1991). *Ptitsy ostrova Sakhalin*. Vladivostok: Vladivostok Institute of Biology and Pedology.

Nekrasov, B.V. (1978). *Ptitsy Volzhsko-Kamskogo Kraja*. Moscow: Nauka.

Newborn, D. (2000). The value of hebridean sheep in controlling invasive purple moor grass. *Aspects of Applied Biology* 58, 191-196.

Newman, J.A., Parsons, A.J., Thornley, J.H.M., Penning, P.D. & Krebs, J.R. (1995). Optimal diet selection by a generalist grazing herbivore. *Functional Ecology* 9, 255-268.

Newton, I. (1998). *Population Limitation in Birds*. Academic Press, London.

Niewold, F.J.J. (1990). The decline of Black Grouse in the Netherlands. In Lumeij, J.T. & Hoogeveen, Y.R. (eds) *The Future of Wild Galliformes in the Netherlands*: 71–81. Amersfoort, Netherlands: Organisatiecommissie Nederlandse Wilde Hoenders.

Niezen, J.H., Charleston, W.A.G., Hodgson, J., Mackay, A.D. & Leathwick, D.M. (1996) Controlling internal parasites in grazing ruminants without recourse to anthelmintics - approaches, experiences and prospects. *International Journal for Parasitology* 26, 983-992.

Niezen, J.H., Robertson, H.A., Waghorn, G.C. & Charleston, W.A.G. (1998) Production, faecal egg counts and worm burdens of ewe lambs which grazed six contrasting forages. *Veterinary Parasitology* 80, 15-27.

Nolan, A.J., Henderson, D.J. & Merrell, B.G. (1995). The vegetation dynamics of wet heaths in relation to sheep grazing intensity. In: *Heaths and Moorland: Cultural*

Lanscapes (eds D.B.A. Thompson, A.J. Hester & M.B. Usher), pp.174-179. HMSO, Edinburgh.

O'Brien, M., Green, R.E. & Wilson, J. (2006). Partial recovery of the population of Corncrakes *Crex crex* in Britain, 1993-2004. *Bird Study* 53, 213-224.

O'Connell, M., Thomas, C.J., Twiss, S.D., Downie, I.S., Evans, P.R. & Whitfield, D.P. (1996). *Functional Ecology of Peatland Animals in the Flow Country of Northern Scotland. I. Habitat Requirements of Breeding Waders Charadrii*. Edinburgh: Research and Advisory services Directorate Report, SNH.

Oglethorpe, D.R. & Sanderson, R.A. (1999). An ecological- economic model for agri-environmental policy analysis. *Ecological Economics* 28, 245-266.

Oglethorpe, D.R., & O'Callaghan, J.R., (1995). Farm level modelling within a river catchment decision support system. *Journal of Environmental Planning and Management* 38, 93-106.

Oglethorpe, D. (2005). Livestock production post CAP reform: implications for the environment. *Animal Science*, **81**, 189-182.

Oksanen, J. (2006). Multivariate analysis of ecological communities in R. Unpublished report.

Onate J.J., Atance I., Bardaji I., Llusia D. (2007) Modelling the effects of alternative CAP policies for the Spanish high-nature value cereal-steppe farming systems. *Agricultural Systems*, **94**, 247-260.

Ormerod, S.J., Barlow, N.D., Marshall, E.J.P. & Kerby, J. (2002). The uptake of applied ecology. *Journal of Applied Ecology* 39, 1-7.

Osoro, K., Benito-Pena, A., Frutos, P., Garcia, U., Ortega-Mora, L.M., Celaya, R. & Ferre, I. (2007). The effect of heather supplementation on gastrointestinal nematode infections and performance in Cashmere and local Celtiberic goats on pasture. *Small Ruminant Research* 67, 184-191.

Osoro, K., Olivan, M., Celaya, R. & Martinez, A. (2000). The effect of *Calluna vulgaris* cover on the performance and intake of ewes grazing hill pastures in northern Spain. *Grass and Forage Science* 55, 300-308.

Osterburg B. & van Horn L. (2006). Assessing the impacts of decoupling EU direct payments from agricultural production and the potential for 're-coupling'. *Outlook in Agriculture*, **35**, 107-113.

Pakeman, R.J. (2004). Consistency of plant species and trait responses to grazing along a productivity gradient: a multi-site analysis. *Journal of Ecology* 92, 893-905.

Pakeman, R.J., Hulme, P.D., Torvell, L. & Fisher, J.M. (2003). Rehabilitation of degraded dry heather *Calluna vulgaris* (L.) Hull moorland by controlled sheep grazing. *Biological Conservation* 114, 389-400.

Palmer, S.C.F. & Hester, A.J. (2000). Predicting spatial variation in heather utilization by sheep and red deer within heather/grass mosaics. *Journal of Applied Ecology* 37, 616-631.

Palmer, S.C.F. (1997). Prediction of the shoot production of heather under grazing in the uplands of Great Britain. *Grass and Forage Science* 52, 408-424.

Palmer, S.C.F., Gordon, I.J., Hester, A.J. & Pakeman, R.J. (2004). Introducing spatial grazing impacts into the prediction of moorland vegetation dynamics. *Landscape Ecology* 19, 817-827.

Palmer, S.C.F., Hester, A.J., Elston, D.A., Gordon, I.J. & Hartley, S.E. (2003). The perils of having tasty neighbours: grazing impacts of large herbivores at vegetation boundaries. *Ecology* 84, 2877-2890.

Park, K.J., Robertson, P.A., Campbell, S.T., Foster, R., Russell, Z.M., Newborn, D. & Hudson, P.J. (2001). The role of invertebrates in the diet, growth and survival of Red Grouse *Lagopus lagopus scoticus* chicks. *Journal of Zoology* 254, 137-145.

Parsons, A.J., Harvey, A. & Johnson, I.R. (1991). Plant animal interactions in a continuously grazed mixture. 2. The role of differences in the physiology of plant-growth and of selective grazing on the performance and stability of species in a mixture. *Journal of Applied Ecology* 28, 635-658.

Partridge, L.W., Trenkel, V.M., Gordon, I.J., Buckland, S.T., Elston, D.A., Milne, J.A., Birch, C.P.D., Forster, J., Mann, A.D. & McLeod, J.E. (1999). HillDeer: The Red Deer Decision Support System - User's Guide. Aberdeen: MLURI/BioSS.

Pearce-Higgins, J.W. & Grant, M.C. (2006). Relationships between bird abundance and the composition and structure of moorland vegetation. *Bird Study* 53, 112–125.

Pearce-Higgins, J.W. & Yalden, D.W. (2003). Variation in the use of pasture by breeding European Golden Plovers *Pluvialis apricaria* in relation to prey availability. *Ibis* 145, 365-381.

Pearce-Higgins, J.W. & Yalden, D.W. (2004). Habitat selection, diet, arthropod availability and growth of a moorland wader: the ecology of European Golden Plover *Pluvialis apricaria* chicks. *Ibis* 146, 335-346.

Pearce-Higgins, J.W. (1999). The Ecology of Golden Plovers *Pluvialis apricaria* in the Peak District. PhD thesis, University of Manchester.

Pearce-Higgins, J.W., Grant, M.C., Robinson M.C., & Haysom S.L. (2007). The role of forest maturation in causing the decline of Black Grouse *Tetrao tetrix*. *Ibis* 149, 143–155.

Penny Anderson Associates (2004). Monitoring of Demonstration Moors in England and Wales – Establishment of Baseline Data. Unpublished interim report to Defra (contract BD1237).

Peresad'ko, L.V. & Koshelev, A.I. (1980). K pitaniyu kulikov na yuge Zapadnoi Sibiri. *Novoe v izuchenii biologii i rasprostraneni kulikov*. Moscow: Nauka.

Peter, D., Young, M.R. & Bentley, C. (2001). The effects of varied grazing management on epigeal spiders, harvestmen and pseudoscorpions of *Nardus stricta* grassland in upland Scotland. *Agriculture Ecosystems & Environment* 86, 39-57.

Picozzi, N. & Hepburn, L.V. (1984). A study of Black Grouse in north east Scotland. *Proceedings of the International Grouse Symposium* 3, 462–480.

Prokof'yeva, I.V. (1980). Pitanie lugovykh vorob'inykh ptits v period gnezdovaniya. *Ornitologiya* 15.

Purvis, O., Coppins, B., Hawksworth, D., James, P. & Moore, D. (1992). The Lichen Flora of Great Britain and Ireland. The Natural History Museum in association with the British Lichen Society, London.

Putwain, P.D. & Rae, P.A.S. (1988). Heathland restoration: a handbook of techniques. British Gas, Southampton.

- Pykala, J. (2005). Plant species responses to cattle grazing in mesic semi-natural grassland. *Agriculture, Ecosystems & Environment* 108, 109-117.
- Pywell, R.F., Webb, N.R., & Putwain, P.D. (1995). A comparison of techniques for restoring heathland on abandoned farmland. *Journal of Applied Ecology* 32, 400-411.
- Ramsay, S.L. & Houston, D.C. (2003). Amino acid composition of some woodland arthropods and its implications for breeding tits and other passerines. *Ibis* 145, 227–232.
- Randin, C., Dirnböck, T., Dullinger, S., Zimmermann, N., Zappa, M. & Guisan, A. (2006). Are niche-based species distribution models transferable in space? *Journal of Biogeography* 33, 1689-1703.
- Ratcliffe, D.A. (1976). Observations on the breeding of Golden Plover in Great Britain. *Bird Study* 23, 63–116.
- Ratcliffe, D.A. (1990). *Bird Life of Mountains and Uplands*. Cambridge University Press, Cambridge.
- Read, J.M., Birch, C.P.D. & Milne, J.A. (2002). HeathMod: a model of the impact of seasonal grazing by sheep on upland heaths dominated by *Calluna vulgaris* (heather). *Biological Conservation* 105, 279-292.
- Redpath, S.M. & Thirgood, S.J. (1997). *Birds of prey and red grouse*. HMSO, London.
- Redpath, S.M. & Thirgood, S.J. (1999) Functional and numerical responses in generalist predators: hen harriers and peregrines on Scottish grouse moors. *Journal of Animal Ecology* 68, 879-893.
- Rice R.W. (1986). Models as a tool in grazing management. In: Gudmundsson O. (ed) *Grazing Research at Northern Latitudes*. Proceedings of a NATO Advanced Research Workshop, Hvanneyri, 1985. pp. 291-299. New York: Plenum Press.
- Robbins, C.T., Hanley, T.A., Hagerman, A.E., Hjeljord, O., Baker, D.L., Schwartz, C.C. & Mautz, W.W. (1987). Role of tannins in defending plants against ruminants: reduction in protein availability. *Ecology* 68, 98-107.

Robertson, P.A., Park, K.J. & Barton, A.F. (2001). Loss of heather *Calluna vulgaris* moorland in the Scottish uplands: the role of Red Grouse *Lagopus lagopus scoticus* management. *Wildlife Biology* 7, 11–16.

Robson, G. (1998). The Breeding Ecology of Curlew *Numenius arquata* on North Pennine Moorland. PhD thesis, University of Sunderland.

Robson, G., Percival, S.M. & Brown, A.F. (2002). The use of marginal farmland by Curlew *Numenius arquata* breeding on upland moors. *Aspects of Applied Biology* 67, 75–84.

Rodríguez, A. & Andrén, H. (1999). A comparison of Eurasian red squirrel distribution in different fragmented landscapes. *Journal of Applied Ecology* 36, 649–662.

Rodwell, J.S. (1991). *British Plant Communities Volume 2: Mires and Heaths*. Cambridge University Press, Cambridge.

Rodwell, J.S. (1992). *British Plant Communities Volume 3: Grasslands and montane communities*. Cambridge University press, Cambridge.

Rook, A.J., Dumont, B., Isselstein, J., Osoro, K., WallisDeVries, M.F., Parente, G. & Mills, J. (2004). Matching type of livestock to desired biodiversity outcomes in pastures – a review. *Biological Conservation* 119, 137-150.

Roper, M. (1999). The outlook for the sheep industry and its impact on sheep research requirements. Invited summaries, Proceedings of the BSAS Annual Meeting, Scarborough.

Roughsedge, T., Amer, P. & Simm, G. (2002). Optimum replacement strategies for suckler cow beef. Conference paper at 7th World Congress on Genetics applied to Livestock Production, Montpellier.

Rural Development Service (2003). *Environmental Cross Compliance: Vegetation change assessment survey, Birkbeck Common, Cumbria*. Unpublished internal report of the National Grazing Management Team (Leeds), Rural Development Service, May 2003.

Rushton, S.P., Ormerod, S.J. & Kerby, G. (2004). New paradigms for modelling species distributions? *Journal of Applied Ecology* 41, 193-200.

Rushton, S.P., Sanderson, R.A., Wildig, J. & Byrne, J.P. (1996). The effects of grazing management on moorland vegetation: a comparison of farm unit, grazing paddock and plot experiments using a community modelling approach. *Aspects of Applied Biology* 44, 211-219.

Sanderson, R.A., Rushton, S.P., Cherrill, A.J. & Byrne, J.P. (1995). Soil, vegetation and space: analysis of their effects on invertebrate communities of a moorland in north-east England. *Journal of Applied Ecology* 32, 506-518.

Sanderson, R.A. & Rushton, S.P. (1995). VEMM - predicting the effects of agricultural management and environmental conditions on semi-natural vegetation. *Computers and Electronics in Agriculture* 12, 237-247.

Sanderson, R.A., Eyre, M.D. & Rushton, S.P. (2005). The influence of stream invertebrate composition at neighbouring sites on local assemblage composition. *Freshwater Biology* 50, 221-231.

SAS Institute (2001). *SAS/STAT User's Guide, Version 8.02*. SAS Institute, Cary NC.

Savory, C.J. (1977). The food of red grouse chicks, *Lagopus lagopus scoticus*. *Ibis* 119, 1-9.

Scandrett, E. (1991). *Regenerating heather moorland*. Macaulay Land Use Research Institute.

Schwartz, C.C. & Ellis, J.E. (1981). Feeding ecology and niche separation in some native and domestic ungulates on the shortgrass prairie. *Journal of Applied Ecology* 18, 343-353.

Senft, R.L., Coughenour, M.B., Bailey, D.W., Rittenhouse, L.R., Sala, O.E. & Swift, D.M. (1987). Large herbivore foraging and ecological hierarchies. *Bioscience* 37, 789-799.

Shepherd, M.J., Anderson, J.M., Bol, R. & Allen, D.K. (2000). Incorporation of ¹⁵N from spiked cattle dung pats into soil under two moorland plant communities. *Rapid Communications in Mass Spectrometry* 14, 1361-1367.

Sibbald A.R., Milne J.A., Mathews K.B. & Birch C.P.D. (2000). Decision support for hill and upland farms in the UK. In: Gagnaux D., Daccord R., Gibon A., Poffet J.R. and Sibbald A.R. (eds) *Livestock Farming Systems – Integrating Animal Science*

Advances into the Search for Sustainability. Proceedings of the European Association for Animal Production 5th International Symposium on Livestock Farming Systems. EAAP Publication No. 97. pp. 284-288. Wageningen: Wageningen Pers.

Siepel, H. (1990). The influence of management on food size in the menu of insectivorous animals. *Proceedings of Experimental and Applied Entomology* 1, 69–74.

Sim, I.M.W., Burfield, I.J., Grant, M.C., Pearce-Higgins, J.W. & Brooke, M. de L. (2007). The role of habitat composition in determining breeding site occupancy in a declining Ring Ouzel *Turdus torquatus* population. *Ibis* 149, 378-385.

Sim, I.M.W., Gregory, R.D., Hancock, M.H. & Brown, A.F. (2005). Recent changes in the abundance of British upland breeding birds. *Bird Study* 52, 261-75.

Simpson, I.A., Kirkpatrick, A.H., Scott, L., Gill, J.P., Hanley, N. & MacDonald, A.J. (1998). Application of a grazing model to predict heather moorland utilization and implications for nature conservation. *Journal of Environmental Management* 54, 215-231.

Sinclair, K.D., Yildiz, S., Quintans, G. & Broadbent, P.J. (1998). Annual energy intake and the performance of beef cows differing in body size and milk potential. *Animal Science* 66, 643-655.

Smith, A.A., Redpath, S.M., Campbell, S.T. & Thirgood, S.J. (2001). Meadow pipits, red grouse and the habitat characteristics of managed grouse moors. *Journal of Applied Ecology* 38, 391-401.

Smith, A.J.E. (2004). *The moss flora of Britain and Ireland*. Cambridge University Press, Cambridge, 1012pp.

Smith, R.S., Charman, D., Rushton, S.P., Sanderson, R.A., Simkin, J.M. & Shiel, R.S. (2003). Vegetation change in an ombrotrophic mire in northern England after excluding sheep. *Applied Vegetation Science* 6, 261-270.

Soil Survey of England & Wales (1983). 1:250 000 Soil Map of England and Wales. Soil Survey of England and Wales, Rothamsted.

- Southwood, T.R.E., Brown, V.K. & Reader, P.M. (1979). The relationships of plant and insect diversities in succession. *Biological Journal of the Linnean Society* 12, 327–348.
- Spalinger, D.E., Hanley, T.A. & Robbins, C.T. (1988). Analysis of the functional response in foraging in the Sitka black-tailed deer. *Ecology* 69, 1166-1175.
- Stace, C.A. (1995). *New flora of the British Isles*. Cambridge University Press, Cambridge, 1226pp.
- Staines, B.W., Balharry, R. & Welch, D. (1995). The impact of red deer and their management on the natural heritage in the uplands. In *Heaths and Moorland - Cultural Landscapes*, eds Thompson, D.B.A., Hester, J.A. & Usher, M.B. Edinburgh: HMSO. 294-308.
- Starling-Westerberg, A. (2001). The habitat use and diet of Black Grouse *Tetrao tetrix* in the Pennine hills of northern England. *Bird Study* 48, 76–89.
- StatSoft, Inc. (2001). *Statistica for Windows [Computer program manual]*. Tulsa, Oklahoma.
- Stephen, L. (2001). *An Investigation into the Dietary Requirements of Curlew *Numenius arquata* Chicks*. BSc thesis, University of Edinburgh.
- Stephens, D.W. & Krebs, J. (1986). The economics of choice: trade-offs and herbivory. *Foraging Theory*. pp. 104-127. Princeton University Press: New Jersey.
- Stillman, R.A. & Brown, A.F. (1994). Population sizes and habitat associations of upland breeding birds in the South Pennines, England. *Biological Conservation* 69, 307-314.
- Stuth, J.W. (1991). Foraging behavior. In: Heitschmidt R.K. and Stuth J.W. (eds). *Grazing management. An ecological perspective*. pp. 65-83. Portland, Oregon: Timber Press.
- Task Force for the Hills (2001). *Report of Findings to the Minister for Agriculture, Fisheries and Food*. London: HMSO.
- Ter Braak, C.J.F. & Šmilauer, P. (1998). *CANOCO reference manual and user's guide to CANOCO for Windows: Software for canonical community ordination (version 4)*. Microcomputer Power, Ithaca, NY.

Ter Braak, C.J.F. & Smilauer, P. (2002). CANOCO reference manual and Cano Draw for Windows User's guide: Software for Canonical Community Ordination (version 4.5) Microcomputer Power, Ithaca, New York, America.

Ter Braak, C.J.F. (1986). Canonical correspondence analysis: a new eigenvector technique for multivariate direct gradient analysis. *Ecology* 67, 1167-1179.

Terry, A.C., Ashmore, M.R., Power, S.A., Allchin, E.A. & Heil, G.W. (2004). Modelling the impacts of atmospheric nitrogen deposition on Calluna-dominated ecosystems in the UK. *Journal of Applied Ecology* 41, 897-909.

Tharme, A.P., Green, R.E., Baines, D., Bainbridge, I.P. & O'Brien, M. (2001). The effect of management for red grouse shooting on the population density of breeding birds on heather dominated moorland. *Journal of Applied Ecology* 38, 439-457.

Theodorou, M.K., Williams, B.A., Dhanoa, M.S., Mcallan, A.B. & France, J. (1994). A simple gas production method using a pressure transducer to determine the fermentation kinetics of ruminant feeds. *Animal Feed Science and Technology* 48, 185-197.

Thirgood, S.J. Redpath, S.M., Haydon, D., Rothery, P., Newton, I. & Hudson, P.J. (2000). Habitat loss and raptor predation: disentangling long and short term causes of red grouse declines. *Proceedings of the Royal Society Series B* 267, 651-656.

Thirgood, S.J., Leckie, F.M. & Redpath, S.M. (1995). Diurnal and seasonal variation in line transect counts of moorland passerines. *Bird Study* 42, 257-259.

Thomas, T.A. (1977). An automated procedure for the determination of soluble carbohydrates in herbage. *Journal of the Science of Food and Agriculture* 28, 639-642.

Thompson, D.B.A., MacDonald, A.J., Marsden, J.H. & Galbraith, C.A. (1995). Upland heather moorland in Great Britain: a review of international importance, vegetation change and some objectives for nature conservation. *Biological Conservation* 71, 163-178.

Tilley, J.M.A. & Terry, R.A. (1963). A two-stage technique for the in vitro digestion of forage crops. *Journal of the British Grassland Society* 18, 104-111.

- Todd, P.A., Phillips, J.D.P., Putwain, P.D. & Marrs, R.H. (2000). Control of *Molinia caerulea* on moorland. *Grass and Forage Science* 55, 181-191.
- Tudor, G. & Mackey, E.C. (1995). Upland land cover change in post war Scotland. In: Usher, M.B., Thompson, D.B.A. (Eds.) *Ecological change in the Uplands*. Blackwell Scientific Publications, Oxford, 28-42.
- Tyler, S.J. & Green, M. (1994). The status and breeding ecology of Ring Ouzels *Turdus torquatus* in Wales with reference to soil acidity. *Welsh Bird Report* 7, 78-89.
- UK Biodiversity Steering Group (1995). *Biodiversity: The UK Steering Group Report, Vol. II: Action plans*. HMSO, London.
- Usher, M.B. & Thompson, D.B.A. (1993). Variation in the upland heathlands of Great Britain: Conservation importance. *Biological Conservation* 66, 69-81.
- van der Wal, R., Pearce, I., Brooker, R., Scott, D., Welch, D. & Woodin, S. (2003). Interplay between nitrogen deposition and grazing causes habitat degradation. *Ecology Letters* 6, 141-146.
- Van Soest, P.J. (1982). *The Nutritional Ecology of the Ruminant*. O & B Books, Oregon.
- Van Soest, P.J., Robertson, J.B. & Lewis, B.A. (1991). Methods for dietary fiber, neutral detergent fibre, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* 74, 3583-3597.
- Vandvik, V., Heegaard, E., Måren, I. E., & Aarrestad, P.A. (2005). Managing heterogeneity: the importance of grazing and environmental variation on post-fire succession in heathlands. *Journal of Applied Ecology* 42, 139-149.
- Vanreusel, W., Maes, D. & van Dyck, H. (2007). Transferability of species distribution models: a functional habitat approach for two regionally threatened butterflies. *Conservation Biology* 21, 201-212.
- Virtanen, R., Edwards, G.R. & Crawley, M.J. (2002). Red deer management and vegetation on the Isle of Rum. *Journal of Applied Ecology* 39, 572-583.
- von Labhardt, A. (1988). Zur nahrung des Braunkehlchens *Saxicola rubetra* wahrend der Jungenaufzucht. *Beih. Veroff. Naturschutz Landschaftspflege Bad.-Wurt.* 51: 179-185.

- Vulink, J.T. & Drost, H.J. (1991). A causal analysis of diet composition in free ranging cattle in reed-dominated vegetation. *Oecologia* 88, 167-171.
- Walker, A.D., Campbell, C.G.B., Heslop, R.E.F., Gauld, J.H., Laing, D., Shipley, B.M. & Wright, G.G. (1982). Soil Survey of Scotland. Macaulay Institute for Soil Research, Aberdeen.
- Wallach, D., Elsen, J.M. & Charpentreau, J.L. (1984). Maintenance energy requirements of grazing sheep: A detailed comparison between models. *Agricultural Systems* 15, 1-22.
- Walton, K.C. (1979). Diet of meadow pipits *Anthus pratensis* on mountain grassland in Snowdonia. *Ibis* 121, 325-334.
- Wang, C.T & Dickerson, G.E. (1991). A deterministic computer simulation model of life-cycle lamb and wool production. *Journal of Animal Science* 69, 4312-4347.
- Welch, D. & Scott, D. (1995). Studies in the grazing of heather moorland in north-east Scotland. VI. 20-year trends in botanical composition. *Journal of Applied Ecology* 32, 596-611.
- Welch, D. (1984). Studies in the grazing of heather moorland in north-east Scotland. II. Response of heather. *Journal of Applied Ecology* 21, 197-207.
- Welch, D. (1985). Studies in the grazing of heather moorland in north-east Scotland. IV. Seed dispersal and plant establishment in dung. *Journal of Applied Ecology* 22, 461-472.
- Welch, D. (1986). Studies in the grazing of heather moorland in north-east Scotland. V. Trends in *Nardus stricta* and other unpalatable graminoids. *Journal of Applied Ecology* 23, 1047-1058.
- Welch, D. (1998). Response of bilberry *Vaccinium myrtillus* L. stands in the Derbyshire Peak District to sheep grazing, and implications for moorland conservation. *Biological Conservation* 83, 155-164.
- Whitby, B. & Wadsworth, R. (1994). A bioeconomic model of heather moorland management and conservation. *Ecological Conservation* 9, 167-177.

Whitby, M. & Grant S.A. (eds) (1990). Modelling heather management, a workshop herald by Department of Agricultural Economics and Food Marketing, University of Newcastle upon Tyne, 46-55.

White, B. & Whitby, M. (1990) .An economic analysis of heather management. In 'Modelling heather management' Whitby, M. and Grant S.A. (eds) Department of Agricultural Economics and Food Marketing, University of Newcastle upon Tyne, 46-55.

Whittingham, M., Wilson, J.D, & Donald, P.F (2003). Do habitat association models have any generality? Predicting skylark *Alauda arvensis* abundance in different regions of southern England. *Ecography* 26, 521-531.

Whittingham, M.J. (1996). Habitat Requirements of Breeding Golden Plover *Pluvialis apricaria*. PhD thesis, University of Sunderland.

Whittingham, M.J., Percival, S.M. & Brown, A.F. (2000). Time budgets and foraging of breeding golden plover *Pluvialis apricaria*. *Journal of Applied Ecology* 37, 632-646.

Whittingham, M.J., Percival, S.M. & Brown, A.F. (2001). Habitat selection by golden plover *Pluvialis apricaria* chicks. *Basic and Applied Ecology* 2, 177–191.

Whittingham, M.J., Stephens, P.A., Bradbury, R.B. & Freckleton R.P. (2006). Why do we still use stepwise modelling in ecology and behaviour? *Journal of Animal Ecology* 75, 1182–1189.

Whittingham, M.J., Swetnam, R.D., Wilson, J.D., Chamberlain, D.E. & Freckleton, R.P. (2005). Habitat selection by yellowhammers *Emberiza citrinella* on lowland farmland at two spatial scales: implications for conservation management. *Journal of Applied Ecology* 42, 270–280.

Wilson, J.D., Morris, A.J., Arroyo, B.E., Clark, S.C. & Bradbury, R.B. (1999). A review of the abundance and diversity of invertebrate and plant food of granivorous birds in northern Europe in relation to agricultural change. *Agriculture, Ecosystems and Environment* 75, 13–30.

Wright, I.A., Jones, J.R., Maxwell, T.J., Russel, A.J.F. & Hunter, E.A. (1994). The effect of genotype X environment interactions on biological efficiency in beef-cows. *Animal Production* 58, 197-207.

Yallop, A.R., Thacker, J.I., Thomas, G., Stephens, M., Clutterbuck, B., Brewere, T., & Sannier, C.A.D. (2006). The extent and intensity of management burning in the English uplands. *Journal of Applied Ecology* 43, 1138-1148.

Yeo, M.J.M. & Blackstock, T.H. (2002). A vegetation analysis of the pastoral landscapes of upland Wales, UK. *Journal of Vegetation Science* 13, 803-816.

Zinov'yev, V.I. (1980). *Ptitsy lesnoi zony Evropeiskoi chasti SSSR rzhankoobraznye*. Kalinin: Kalinin State University Press.